

Effects of Gaseous Ozone (O₃) on Artificial Population and Aflatoxin B₁ Production by *Aspergillus flavus* in Stored Whole Cinnamon in Iraqi Markets

Diyaa Aljaza

Asis Prof., Department of pathological Analyses/ College of Science, University of Thi-Qar, Iraq

Abstract

Stored whole cinnamon can become contaminated with aflatoxins. For public health requirements cinnamon should be produced free of AFB₁ or below EU legislation limit (5 µg/kg), so in this study, examined the efficacy of different concentrations (0, 300 and 600 ppm) of ozone on fungal populations (CFU) and AFB₁ production of *A. flavus* on stored whole cinnamon at three a_w levels (0.92, 0.94 & 0.96 a_w) at 25°C and exposure time (60 min). In general, the results presented in this study showed that the populations AFB₁ produced of *A. flavus* in artificially contaminated whole cinnamon were significantly affected by O₃ treatment.

Keywords: Cinnamon; Ozone; Population; Aflatoxin B₁; *Aspergillus flavus*.

Introduction

Cinnamon produced in tropical countries and used as ingredients in traditional foods as food supplements or medicines⁽³⁾. Cinnamon can be contaminated with AFB₁ during pre or post-harvest, processing, handling and storage⁽²⁰⁾. Aflatoxins (AFs) are mycotoxins produced by many fungal species, especially *Aspergillus flavus*, *A. nomius* and *A. parasiticus*. These AFs are carcinogenic, teratogenic and mutagenic in human and animals. There are many types of AFs one of them Aflatoxin b₁ (AFB₁), it is the most toxic because classified by International Agency for Research on Cancer (IARC) as class 1a carcinogenic in human and animals⁽¹²⁾.

Ozone (O₃) is a strong germicide and has three oxygen atoms. It is used in food industries as an antifungal agent prevents fungal growth in raw materials and used as a preservative in packaging foods⁽²³⁾. O₃ is produced naturally as a colourless or bluish gas by

two methods and artificially by three methods. The most common method for O₃ production is a corona-discharge method (CDM)^(16;9). It will be considered in this study to control fungal population and prevent contamination of cinnamon with AFs.

Natural levels of O₃ are between 0.01-0.15 ppm but exposure to high levels produce some harmful effects in health such as respiratory diseases⁽¹¹⁾ and cardiovascular diseases⁽⁶⁾. Federal occupational safety and health administration (OSHA) recommended a legislative limit for exposure to O₃ of 0.1 ppm for 8 hr continuously or 0.3 ppm for 15 min⁽²²⁾.

The efficacy of O₃ on fungal species is depending on fungal tolerance to exposure, O₃ (concentration), exposure time, spore morphology and moisture content of substrate⁽¹⁾. Moreover, O₃ treatment can cause negative impacts in the quality of the food such as oxidation of lipids, change colour, modified of some vitamins and phenolic compounds^(18;7).

The effect of O₃ on colonization and AFB₁ production of *A. flavus* in peanuts, wheat, chillies, Brazil nut, corn, maize and red peppers have been studied previously,^(21;5;2;19;4;17;10;13). To my knowledge, there very few studies on cinnamon. The objective of this

Corresponding Author:

Asis Prof. Dr Diyaa Aljaza

Department of pathological Analyses, College of Science, University of Thi-Qar, Iraq.

Email address: diyaamt71@gmail.com

study was to examine (a) the efficacy of O₃ on the total fungal colonization and AFB₁ production of *A. flavus* in artificially contamination on whole dried cinnamon stored at 25°C stored for 20 days.

Material and Methods

Chemicals and reagents

The AflaStar™ – Immunoaffinity Column (AIC) were purchased from Romer Labs PLC (Tulln, Venna, Austria). Aflatoxins Stock was purchased from (Romer Lab UK Ltd, Cheshire, UK). Mealt Extract Agar was purchase from (Oxoid, Basingstoke, Hampshire, UK). Methanol and Acetonitrile were obtained from (Fisher Scientific, UK). All solvents were of HPLC grade.

Collection of cinnamon samples

300 g of whole dried cinnamon were randomly collected from Iraqimarkets and direct plated on MEA and then incubated at 25°C for seven days for fungal isolation.

Preparation of fungal conidial inoculums

Spore inoculums was made from *A. flavus* (DJC₁) isolated from cinnamon, these spores grown on MEA for 7 days and removed by sterile loop and then flooded with 9 ml of sterile water containing 0.025% (w/v) Tween 80.

Media preparation

Malt extract agar (MEA)

This medium was used for culturing, isolating and enumerating fungal species. (15 g MEA powder and 300 ml distilled water (D.W). A small amount of antibiotic agent, chloramphenicol (0.05 g/L) was mixed with the components to prevent bacterial growth. After autoclaved for 1 hr at 121°C the sterile medium was cooled and poured into 9 cm serial Petri plates and left to cool, and kept at 4°C in polyethene bags until used.

Preparation of Adsorption curve

A moisture absorption curve was designed to determine the quantities of water required to add to cinnamon samples to modify the moisture contents to the treatment target a_w values. This curve was prepared by adding different amounts of distilled water to whole cinnamon and equilibrating at 4°C for 24 hrs. The a_w was

measured with an Aqua Lab 4TE.

Preparation of chilli samples

Thirty grams of whole dried cinnamon were weighed into 9 solid culture vessels and then stoppered with plastic lids with a permeable membrane. The samples divided into three levels of a_w conditions (0.92, 0.94 and 0.96a_w). Each group of chillies adapted by adding the appropriate volume of distilled water based on the moisture absorption curve to obtain the required target levels. All samples were shaken on a (KS 501 digital orbital shaker) for 1hr and then kept at 4°C for 24 hrs to allow absorption and equilibration. Each vessel was inoculated with a spore suspension of *A. flavus* (10² spores/ml) by add 0.25 ml of spore solution to each vessel. All samples mixed well and exposed to a different concentration of O₃ (0, 300 and 600) ppm for 60 min at (6L/min) in 50 ml class tube system. Controls were prepared by exposure to air only. Each treatment divided into three separate replicates at each a_w levels and stored in plastic chambers at 25°C for 20 days. Each plastic chamber also contained 500 ml beaker containing a glycerol/water solution of the same a_w to maintain the ERH. The *A. flavus* populations present initially and after 10 and 20 days were quantified at each treatment. The samples were stored at -20°C for AFB₁ analyses at each treatment condition.

Enumeration of *A. flavus* populations on stored whole cinnamon

One g of cinnamon from each treatment and replicate was weighed and mixed with 9 ml of sterile D.W with 0.025% (v/v) Tween 80 to obtain the 10⁻¹ dilution. All samples were then serially diluted after vigorous shaking at each dilution using a vortex mixer. 100 µl of each dilution was spread plate onto MEA plates modified a_w at the same a_w of storage a_w. All plates were incubated at 25°C for seven days the colonies growing were then enumerated.

Extraction of AFB₁ from chillies using HPLC

Weighted three g of cinnamon from each treatment and replicates and placed in 30 ml tube. Added 12 ml of extraction solution (**acetonitrile / water**) (6/4 (v/v)) at 15-18°C. Then the samples have been shaken for 1 hr. The extracts were filtered into contenier using a

funnel containing a Whatman filter paper (12.5 cm). The extracts were diluted with PBS (Phosphate buffered saline) (pH 7.4) until the content of acetonitrile is lower than 5% (v/v) and (7 pH). The diluted extracts were transferred to (Romer, IAC) and allowed to pass through the column using a syringe barrel as a reservoir. The flow rate should be not exceeding 1-3 ml/min. The column and the extract must be at 15-18°C. The IAC should be rinsed with 2 x 10 ml of distilled water. Added 2 ml of methanol directly to the column and calculated the elution in a new Eppendorf tube 2 ml. The elution was evaporated under reduced pressure at 40°C using a rotary evaporator. Added 1 ml of water: methanol (0.5:0.5) into toxin and filtered the extracts through Nylon 13 mm, 0.2 µm filter directly into amber HPLC vial and then stored at -20°C until analyses using HPLC.

Statistical Analysis

The data obtained from this study were determined by using the Shapiro Wallis Test for (Non-normality)

and one-way ANOVA for (Normality). The significant statistical level was at $P < 0.05$ for both factors and their interactions.

Results

The effects of O₃ (ppm) on artificially populations of *A. flavus* on whole cinnamon stored at different a_w levels.

Figure 1 shows the effect of exposure to different doses of O₃ on the colonisation by *A. flavus* on stored whole cinnamon for 20 days at 25°C and different a_w levels. The results showed that there were clear effects of incubation time and a_w on the populations in stored whole cinnamon and significant effects of O₃ concentrations except on initial colonization at first day of incubation especially, at 0.92 and 0.96 a_w. Statistically, the populations were significantly affected by O₃ concentrations, time, ppm x time, ppm x a_w and a_w but unaffected by time x a_w and ppm x time x a_w (Table 1).




Figure 1 (a,b &c): Effect of O₃ (ppm) on populations of *A. flavus* (DAJ₁) isolated from stored whole cinnamon at 25°C at (0.92, 0.94 and 0.96a_w) over 20 days. Bars indicate SDM.

The effects of O₃ (ppm) on control the AFB₁ contamination of *A. flavus* in artificially contamination on stored cinnamon at different a_w levels.

Figure 2 showed the efficacy of O₃ concentrations on AFB₁ contamination in stored whole cinnamon for 20 days with four a_w levels at 25°C and exposure time was 60 min. The data of this study reported that AFB₁ of *A. flavus* gradually decreased when increased O₃ concentrations except at 10 days at 0.92 a_w. Also, the AFB₁ production affected by a_w and time. Table 1 shows that there were significant effects of O₃ ppm, a_w, time, ppm x a_w, ppm x time and a_w x time but unaffected by ppm x a_w x time.




Figure 2 (a,b &c): Efficacy of O₃ (ppm) on AFB₁ contamination of *A. flavus* isolated from stored cinnamon at 25°C at (0.92, 0.94 and 0.96a_w) over 20 days. Bars indicate SDM.

Table 1: Summarise statistical analyses of effects of O₃ (ppm) on total populations and AFB₁ contamination of *A. flavus* on whole cinnamon at 25°C under three a_w levels (0.92, 0.94 and 0.96a_w) over 20 days storage. ANOVA test (normality data) and Shapiro Wills test (non-normality data).

Factors	Effects	Actions
ppm	S ^b	Population
	S	AFB ₁ Production
incubation time	S ^b	Population
	S	AFB ₁ Production
a _w	S ^b	Population
	S	AFB ₁ Production
ppm x time	S ^a	Population
	S	AFB ₁ Production
ppm x a _w	S ^a	Population
	S	AFB ₁ Production
ppm x time x a _w	NS ^a	Population
	NS	AFB ₁ Production
time x a _w	S ^a	Population
	S	AFB ₁ Production

NS, not significant S, significant a) ANOVA test b) Shapiro Wills test

Discussion

In this study different gaseous O₃ concentrations (0, 300 and 600 ppm; flow rate of 6 L/min) were used in order to know the effect of O₃ treatment on fungal populations and AFB₁ production in stored cinnamon inoculated with *A. flavus* (DJC₁) 10² spores/ml at three a_w levels (0.92, 0.94 and 0.96 a_w) for 20 days. The fungal population and AFB₁ production has been measured at (T=0, 10 and 20 days) and compared with control (exposure air only). The results showed that the fungal population affected by gaseous O₃ treatment at all a_w levels and there clear impact of ozonation on AFB₁ content in whole cinnamon (artificial contamination). To my knowledge, no previous studies conducted on the effects of O₃ exposure on fungal and AFB₁ contamination in stored cinnamon.

However, there are some studies done on the efficacy of O₃ concentrations on fungal populations and mycotoxins production in some products. For example, ⁽¹⁾ have shown that the populations of three ochratoxigenic strains were significantly reduced in the coffee beans after exposure to 600 ppm O₃ at 0.90 and 0.95 a_w compared with control. This data were similar to the findings in the current study. Similar to Akbar⁽²²⁾ presented that the fungal populations of *A. flavus* in peanut samples inoculated with (10⁵ spores/g) were reduced when exposed to high concentrations of O₃. ⁽⁸⁾ found that the fungal contamination in Brazil nuts was completely stopped from the day of the application when exposure to 31 ppm O₃ for 5 hrs. ⁽¹⁵⁾ suggested that the fungal contamination of *A. parasiticus* in maize will be inhibition in a 63% after treatment of kernel surface with 50 ppm O₃.

For AFB₁ contamination, the findings agree with other studies, for example, ⁽⁵⁾ found that AFB₁ content was degraded in artificially contaminated wheat after exposure to 20 and 400 ppm O₃ for 5, 10, 15 and 20 min. ⁽²²⁾ found that AFB₁ produced in artificially contaminated peanuts was reduced after exposed to 100, 200 and 400 ppm O₃. Last studies by ⁽¹⁾ presented that the OTA content in artificially contaminated coffee stored at 0.75, 0.90 and 0.95 a_w and 30°C for 12 days was significantly reduced when exposure to 600 ppm O₃ for 1 hr. Recent studies on red pepper by ⁽¹⁴⁾ showed that AFB₁ contents in artificially contaminated red pepper treated with O₃ at

three doses 20 mg/L, 40 mg/L and 80 mg/L for 40 min were reduced depend on O₃ doses.

Conclusion

This study showed that *in situ* the populations and AFB₁ production in stored whole cinnamon inoculated with *A. flavus* spores at different a_w levels were relatively affected by O₃ treatment.

Ethical Clearance: The Research Ethical Committee at scientific research by ethical approval of both MOH and MOHSE in Iraq

Conflict of Interest: Non

Funding: Self-funding

References

- 1) **Akbar AH and Magan N.** Growth and Ochratoxin production by *Aspergillus* species in Coffee beans: Impact of climate change and control using O₃ (PhD thesis). Canfield University, Canfield, Beds., MK43 0AL, U.K. 2015.
- 2) **Aljaza D, Magan N.** Mycotoxins in chillies. Diversity of mycotoxigenic fungi, ecology and control strategies using food grade preservatives and gaseous ozone (PhD thesis). Canfield University, Canfield, Beds., MK43 0AL, U.K. 2019.
- 3) **Al-juraifani AA.** Natural occurrence of fungi and aflatoxins of cinnamon in the Saudi Arabia. *African Journal of Food Science*. 2011;8: 460-65.
- 4) **Chen R, Ma F, Li P, Zhang W, Ding X, Zhang Q, Li M, Wang Y and Xu B.** Effect of ozone on aflatoxins detoxification and nutritional quality of peanuts. *Food Chemistry*. 2014; 146: 284-288.
- 5) **El-Desouky TA, Sharoba AMA, Al El-Desouky, El-Mansy HA and Khayria Naguib.** Effect of Ozone Gas on Degradation of Aflatoxin B₁ and *Aspergillus flavus* Fungal. *Journal Environment Analytic Toxicology*. 2012; 2: 2161.
- 6) **Frampton MW, Pietropaoli A, Dentler M, Chalupa D, Little EL, Stewart J, Frasier L, Oakes D, Wiltshire J, Vora R. and Utell MJ.** Cardiovascular Effects of Ozone in Healthy Subjects with and without Deletion of Glutathione-Transferees M₁. *Inhal Toxicology*. 2015; 27: 113-119.

- 7) Gabler FM, Smilanick JL, Mansour MF and Karaca H. Influence of fumigation with high concentrations of ozone gas on postharvest greymould and fungicide residues on table grapes. *Postharvest Biology and Technology*. 2010; 55(2): 85-90.
- 8) Giordano BNE, Simao Manfio D, Galvao S, Scussel JN, Scussel VM. Reduction of in-shell Brazil nut (*Bertholletia excels H. B. K.*) aflatoxin contamination by ozone gas application during storage. *Julius- Kuhn-Archives*. 2010; 425: 574-580.
- 9) Goncalves AA. Ozone-an emerging technology for the seafood industry. *Brazilian Archives Biology and Technology*. 2009; 52: 1527-1539.
- 10) Halima ZH, Riyam HT and Amna MA. Study the effect of ozone gas and ultraviolet radiation and microwave on the degradation of aflatoxin B₁ produce by *Aspergillus flavus* on stored Maize grains. *Journal of Agriculture and Veterinary Science*. 2015; 8: 5-12.
- 11) Horstman DH, Ball BA, Brown J, Gerrity T& Folinsbee LJ. Comparison of pulmonary responses of asthmatic and no asthmatic subjects performing the light exercise while exposed to a low level of ozone. *Toxicology and Industrial Health*. 1995; 11: 369-385.
- 12) International Agency For Research on Cancer (IARC). Agents Classified by the IARC Monographs. Vol. Le 104. International Agency for Research on Cancer. Available online at <http://monographs.Iarc.fr/ENG/Classification/index.phpresearch>. 2012; 10: 100-78.
- 13) Inan F, Pala M and Doymaz I. Use of ozone in detoxification of aflatoxin B₁ in red pepper. *Journal of Stored Products Research*. 2007; 43: 425-429.
- 14) KamberU, Gulbaz G, Aksu P and Dogan A. Detoxification of aflatoxin B₁ in red pepper (*Capsicum Annum l.*) by ozone treatment and its effect on microbiological and sensory quality. *Journal of Food Processing and Preservation*. 2017; 41: e13102.
- 15) Kell SA, Mason LJ, Maier, D. E. and Woloshuk, C. P., 2001. Efficacy and fumigation characteristics of ozone in stored maize. *Journal Stored Products Research* 37, 371-382.
- 16) Kim, J. G., Yousef, AE & Chism GM. Use of ozone to inactivate microorganisms on lettuce [Cited by Xu, L., 1999. Use of ozone to improve the safety of fresh fruits and vegetables]. *Food Technology*. 1999;53: 58-61.
- 17) Luo X, Wang R, Wang L, Li Y, Bian Y and Chen Z. Effect of ozone treatment on aflatoxin B₁ and safety evaluation of ozonized corn. *Journal of Food Control*. 2014a; 37: 171-176.
- 18) Patil S, Torres B, Tiwari BK, Wijngaard HH, Bourke P, Cullen PJ and Valdramidis VP. Safety and quality assessment during the ozonation of cloudy apple juice. *Journal of Food Science*. 2010; 75(7): 437-43.
- 19) Sahab A, Hassanien F, El-Nemr S, Abdel-Alim H and Abdel-Wahhab M. Effect of ozone gaseous on aflatoxin degradation and fat and protein content in peanut seeds. *Journal of Applied Sciences Researches*. 2013; 9: 2170-2175.
- 20) Skrijar MM, Jankovic VV, Moracanin SMV, Vukojevic JB. Xerophilic moulds isolated from spices used in meat industry as potential producers of mycotoxins. In *Proceedings National Science Matika Srpska Novi Sad*. 2012; 123: 7-16.
- 21) Sultan, Y. Biodiversity of Mycotoxigenic *Aspergillus* Species in Egyptian Peanuts and Strategies for Minimizing Aflatoxin Contamination (PhD thesis). Canfield University, Canfield, Beds., MK43 0AL, U.K. 2010.
- 22) Suslow, TV. Ozone applications for postharvest disinfection of edible horticultural crops. *ANS Publication*. 2004; 8133: 1-8.
- 23) US FDA. United States Food and Drug Administration, Code of Federal Regulations 21 CFR part. 2001; 173: 368.