

Prevalence and Correlates of Vitamin D Inadequacy in a Sample of Iraqi People

Ismael Hasan Jawad¹, Hasan ALwan Baiee², Issam Salim Ismail³

¹ Ph.D. Student: University of Babylon College of Nursing , ² Prof. Dr. University of Babylon- Hammurabi College of Medicine, ³ MPH, Public health specialist: Wasit Directorate of Health

Abstract

Background: Vitamin D had been identified as a common metabolic/endocrine abnormality. Middle East countries had a very high rate of vitamin D inadequacy, vitamin D deficiency was associated with non-communicable and infectious diseases.

Objective : To identify the prevalence of Vitamin D level and its correlates among Iraqi people in Al-Hilla city-Babylon province

Methods : A cross-sectional study was conducted on a convenient sample of individuals from all age groups and both genders who attended eight primary health care centers and individuals in Al-Hilla city\Babylon province\Iraq, the study period was started from May through October , 2019. A pretested questionnaire was used for data collection , the questionnaire included information regarding socio demographic characteristics of participants and their history of chronic diseases. The following measurements were done: level of serum vitamin D , blood pressure, blood sugar , body weight and weight and the serum of vitamin D .

Results: The sample included 188 participants their mean age was 64.8 ± 17.25 years. The female to male ratio 1.7:1, More than half of participants (56%) resident in urban areas . Vitamin D mean level was $(20.7 \text{ ng/mL} \pm 14.9 \text{ SD})$. The vast majority (80%) of the sample have vitamin D inadequacy in which (66% have deficiency and 14% have insufficiency), while 37 (20%) of the sample have vitamin D adequacy.

A significant inverse relationship between vitamin D serum level and hypertension, type two diabetes mellitus, increase body weight.

Conclusion: there was a high prevalence of vitamin D inadequacy among Iraqi people mainly in females, a significant association between low vitamin D level and hypertension, type two diabetes mellitus, and obesity.

Keywords: Prevalence, Vitamin D inadequacy, associated factors, Iraq

Introduction

Vitamin D inadequacy is a high priority public health problem globally and affected all ages. Vitamin D inadequacy (both deficiency and insufficiency) was a problem even in sunny countries. This problem

particularly was high in the Middle East, especially among females, it had been estimated that 20–100% of USA, Canadian and European elderly men and women were vitamin D deficient. Additionally, other age groups such as; children, young adults and the middle aged are also at risk of vitamin D deficiency. ^{1,2} Prevalence of Vitamin D deficiency was common among elders nursing home residents ³ Low levels of vitamin D have been implicated in a wide variety of conditions and highly prevalent in the geriatric population, including fractures, functional limitations, cancer, cardiovascular disease.

Corresponding author:

Hasan ALwan Baiee/ University of Babylon- Hammurabi College of Medicine.
email: hassanbaiee@yahoo .com

Evidence suggests that vitamin D supplementation may decrease mortality rates⁴. The prevalence of vitamin D inadequacy varied not only between different countries, but there were intra country variations. It may be as low as 3%⁵ and 11.3%⁶ or as high as 87.8% and 98%^{8,9}. Information from different studies on the vitamin D serum levels among Iraqis over the last years indicate a resurgence of vitamin D inadequacy in otherwise healthy and non-healthy members both males and females in all age groups of the population¹⁰⁻²¹. An inverse relationship was identified between vitamin D serum levels and a number of chronic disease-predisposing conditions such as metabolic syndrome insulin resistance, hypertension, and obesity²²⁻²⁵. This suggests that vitamin D insufficiency/deficiency may play a role at the early stages of the natural history of many chronic diseases^{26,27}. This study was done to measure the prevalence of vitamin D inadequacy among Iraqi population and to identify the associations between vitamin D levels and certain chronic diseases such as hypertension, type two diabetes mellitus, and obesity or overweight.

Methodology

This is across sectional observational descriptive study which includes a convenient sample of apparently healthy individuals from all age groups and both genders living in Al- Hilla City, Babylon province, Iraq, the period of the study started from the first of May through September 2019, the sample size was calculated according to the sample size calculation equation with 95% confidence level, 188 individuals who are the attendees of eight randomly selected primary health care centers and the residents of Babylon elders home are participated voluntarily in this study after obtaining their informed oral consents, after explaining the objectives and the measurements procedures of the study. This study was approved by the Ethics Committee of College of nursing - University of Babylon. A pretested questionnaire (contained questions about demographic characteristics of the participants, their past medical history and history of individual life style habits), this tool was used to interview the participants. Measurements of mean systolic and diastolic blood pressure of each participant were done single handy by one of the researchers. Blood pressure was measured in the sitting position. Patients were kept seated quietly for at least 5 min in a chair. Blood pressure was measured

using a sphygmomanometer with an appropriate cuff the cutoff values, blood pressure more than 140/90 considered hypertension or those who diagnosed previously as hypertensive and taking antihypertensive drugs. Peripheral venous blood samples were collected from each participant to measure Serum level of Vitamin D which was measured by chemo immunoassay method (maglumi instrument) and the random blood sugar, blood sugar level above 200mg/dL considered diabetic as well as individual who was previously diagnosed as type 2 diabetes mellitus. Vitamin D levels are divided into three categories .deficient < 20ng/ml, insufficient between 23-29ng/ml and the normal level is 30-100 ng/ml^{28,29}. body weight and height were measured to calculate body mass index according to the WHO classification³⁰.

Data were analyzed by using (spss) version (23). Continuous quantitative data were expressed as mean and \pm SD, while non-parametric numerical data were expressed as median, with the value of categorical data presented as a number (percentage), t student test and analysis of variance were calculated. Chi square test was done and the Kruskal–Wallis and Mann–Whitney tests were used to compare non-parametric and quantitative variables between groups. P-value < 0.05 considered statistically significant.

Results

A sample of 188 individuals from different settings were participated their mean age \pm SD was 64.8 ± 17.25 years ranged from 3 to 89 years; older people ≥ 60 years formed 83% of them as shown in figure [1]. Female to male ratio 1.7:1, figure [2]. More than half of participants (56%) resident in urban, figure [3]. Vitamin D mean level was (20.7 ng/mL \pm 14.9 SD). The vast majority 151 (80%) of the sample have vitamin D inadequacy in which (66% have deficiency with level ≤ 20 ng/mL and 14% have insufficiency with level 21-29 ng/mL), while 37 (20%) of the sample have vitamin D adequacy with level ≥ 30 ng/mL, figure [4].

Table 1: shows the mean rank differences for vitamin D level according to the demographic characteristics depending on Mann- Whitney ranking statistical test, women have asinificant low vitamin D level, vitamin D inadequacy increased significantly with advancing age while no significant difference was detected in place of residency.

Mean \pm SD differences of blood pressure, blood sugar and BMI among participants with vitamin D inadequacy. Low vitamin D is associated significantly and inversely with blood pressure both systolic and diastolic together with high random blood sugar and the Body Mass Index table(2). Table (3) and figures 6,7,8 and figure 9 reveal the inverse correlation between serum vitamin D and blood pressure (systolic, diastolic), blood sugar and the Body Mass Index respectively while there is a positive correlation between vitamin serum level and age (figure 5).




Figure 1: Frequency distribution of sample according to age groups (N=188).




Figure 2: Gender distribution (N=188) Figure 3: Urban-rural distribution (N=188)




Figure 4: Sample distribution based on vitamin D adequacy (N=188)

Table 1: Mean rank differences for vitamin D level according to the demographic characteristics (N=188).

Demographic ,		Vitamin D level	Mann-Whitney U	Sig.
Gender	male (n=69)	106.51	3277.000	0.021
	female (n=119)	87.54		
Age groups (years)	≤ 59 (n=32)	74.69	1862.000	0.024
	> 60 (n=155)	98.56		
Residency	urban	93.47	4236.500	0.767
	rural	95.84		

Table 2: Mean ± SD differences of blood pressure, blood sugar and BMI among participants with vitamin D sufficiency, insufficiency and deficiency (N=188).

Measurements ,	Vitamin D levels			F*	Sig.
	Sufficiency (n=37)	Insufficiency (n=27)	Deficiency (n=124)		
Systolic BP (mmHg)	130 ± 10.7	138 ± 14.6	144 ± 16.2	13.44	0.0001
Diastolic BP (mmHg)	84.0 ± 5.7	86.0 ± 7.7	90.5 ± 9.1	9.83	0.0001
blood sugar level (mg/dL)	135 ± 47.6	176 ± 53.6	216 ± 64.7	26.97	0.0001
BMI	26.8 ± 4.8	30.3 ± 7.3	35.1 ± 8.8	15.90	0.0001

* ANOVA Test

Table 3: Correlation of Vitamin D level with age, systolic/ diastolic blood pressure, blood sugar and BMI (N=188)

Correlation Coefficient*		Variables ,				
		Age	Systolic	Diastolic	B. sugar	BMI
Vit. D	r	0.156	-0.386-	-0.384-	-0.429-	-0.348-
	Sig.	0.033	0.0001	0.0001	0.0001	0.0001
Age	r		0.054	0.026	-0.014-	-0.200-
	Sig.		0.461	0.726	0.848	0.010
Systolic	r			0.792	0.555	0.290
	Sig.			0.0001	0.0001	0.0001
Diastolic	r				0.515	0.231
	Sig.				0.0001	0.0001
B. sugar	r					0.350
	Sig.					0.0001

* Correlation is significant at the 0.05 level (2-tailed).




Figure 5: Correlation between age (years) and vitamin D level (ng/mL), (N=188).




Figure 6: Correlation between systolic blood pressure (mm Hg) and vitamin D level (ng/mL), (N=188).




Figure 7: Correlation between diastolic blood pressure (mm Hg) and vitamin D level (ng/mL), (N=188).




Figure 8: Correlation between blood sugar level (mg/dL) and vitamin D level (ng/mL), (N=188).




Figure 9: Correlation between BMI and vitamin D level (ng/mL), (N=164).

Discussion

This study reveals that there is alarming high prevalence(80%) of vitamin D inadequacy in a sample of Iraqi people, vitamin D inadequacy varied between

different countries, It may be as low as 3% ⁵ and 11.3% in Irbid in Jordan [6] or as high as 87.8% among healthy Saudi Arabian men ⁷. About 90% of the Qatari population may have insufficient levels of serum vitamin

D²⁷. in the United States, approximately 25% were at risk of vitamin D insufficiency and 8% were deficient¹⁷. Our estimate is similar, if not higher, than that reported from other Arab and Gulf countries like United Arab Emirate³². Prevalence among adolescent Saudi girls and women indicates the severity of the insufficiency with up to 80%³³. This high prevalence of low vitamin D status is thought to be linked to socio-cultural factors, the habit of not fortifying dairy products, and poor vitamin D supplementation³²⁻³⁵. The prevalence of vitamin D inadequacy in this study is slightly less than what was reported in recent study among females in Karbala Province – Iraq¹⁴ but it is a bit higher than what was reported by other local study conducted on women at their reproductive age¹⁸. The finding of the current study depicts an inverse correlation between low serum vitamin D and both systolic and diastolic blood pressure, random blood sugar and obesity this findings are similar to the findings of other studies^{15,16,26,27}.

In conclusions alarming high prevalence of vitamin D deficiency among Iraqi people specially among females have been demonstrated, a positive significant associations between vitamin D inadequacy and hypertension, type 2 diabetes mellitus and obesity, we suggested the carrying out of large scale, studies that might lead to effective reforming actions by policy makers in Iraq regarding vitamin D food fortification and the establishment of interventional preventive local program. Health professionals, policy makers, and the general public in Iraq should be aware of the high prevalence of vitamin D deficiency and the associated health risks.

Acknowledgement: We thank all participants for their cooperation.

Financial Disclosure: There is no financial disclosure.

Conflict of Interest: None to declare.

Ethical Clearance: All experimental protocols were approved under the University of Babylon and all experiments were carried out in accordance with approved guidelines.

References

1. Palacios C, Gonzalez L. Is vitamin D deficiency a major global public health problem?. *J Steroid Biochem Mol Biol*, (2013), <http://dx.doi.org/10.1016/j.jsbmb.2013.11.003>
2. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline [published correction appears in *J Clin Endocrinol Metab*. 2011 Dec;96(12):3908]. *J Clin Endocrinol Metab*. 2011;96(7):1911-1930.
3. Arnljots, R., Thorn, J., Elm, M. *et al.* Vitamin D deficiency was common among nursing home residents and associated with dementia: a cross sectional study of 545 Swedish nursing home residents. *BMC Geriatr* 17, 229 (2017). <https://doi.org/10.1186/s12877-017-0622>
4. Bode LE, PharmD, Mallory Brown X MM Brown MM. Vitamin D Supplementation for Extraskelatal Indications in Older Persons. 2020, 21(2):164-1
5. Al-Jefree NM, Lee P, Ahmed F. Knowledge and attitudes about vitamin D, and behaviors related to vitamin D in adults with and without coronary heart disease in Saudi Arabia. *BMC Public Health*. 2017; 17, 266. doi:10.1186/s12889-017-4183-1.
6. Abdul-Razzak KK, Ajlony MJ, Khoursheed AM, Obeidat BA. Vitamin D deficiency among healthy infants and toddlers: a prospective study from Irbid, Jordan. *Pediatr. Int*. 2011; 53, 839-845.
7. Ardawi MS, Sibiany AM, Bakhsh TM, Qari MH, Maimani AA. High prevalence of vitamin D deficiency among healthy Saudi Arabian men: relationship to bone mineral density, parathyroid hormone, bone turnover markers, and lifestyle factors. *Osteoporos Int*. 2012; 23, 675-686.
8. El-shafie DE, Al-Khashan HI, Mishriky AM. Comparison of vitamin D deficiency in Saudi married couples. *Eur J Clin Nutr*. 2012; 66, 742-745
9. Al-Elq AH. The status of Vitamin D in medical students in the preclerkship years of a Saudi medical school. *J Family Community Med*. 2012; 19, 100-104
10. Jaff P, Rasheed B, Faraj K, Rachid S. Vitamin D Status Among Women Living in Sulaimani, Kurdistan Region and HPLC-MS Analysis for Measuring Vitamin D in Organic and Non-organic Eggs, Kirkuk University Journal /Scientific Studies (KUJSS); 2017; 12: 1992– 0849.
11. About SF, Al-Tu'ma FM, Al-Karem LA.

- Deficiency of Vitamin D and Iron in Anemic Female Iraqi Patients; Human Journals; 2017; 8: 87-96.
12. Al-Rubaye F. Serum Concentration of Vitamin D in Preeclampsia, the Iraqi Postgraduate Medical Journal; 2011; 10: 220-3
 13. Althuwaini T, AL-Thweeni S, AL-Shibany Y. Low Maternal Serum Vitamin D Level at Early Pregnancy and Subsequent Risk of Gestational Diabetes in Women with History of Previous Gestational Diabetes, Karbala J. Med.; 2016; 9: 2543-9.
 14. Al Janaby S, Al Timimy, Al yassery R. Prevalence of Vitamin D Deficiency of Females in Karbala, Iraq, 2017. Karbala J. Med. Vol.13, No.1, June, 2020:2319-25.
 15. Hamza PJ, Hasan MK. The association between Vitamin D level and body mass index in a sample of childbearing age women in Erbil. Med J Babylon 2020;17:165-71.
 16. Abdulrahman RM, Rahman BMA. Prevalence of Vitamin D level in the serum of patients living in Erbil city, Iraq, referred to private clinical laboratory and effect of age and sex on it. J Biol Res Bollet Della Soc Italiana Biol Sperimentale 2018;91:8-11.
 17. Jawad IH, Baiee HA. Prevalence of Vitamin D deficiency and its correlate with overweight and obesity in community-dwelling old adults. Med J Babylon 2020;17:36-40.
 18. Hantoosh HA, Mahdi MH, Imran BW, Yahya AA. Prevalence of Vitamin D deficiency in Iraqi female at reproductive age. Med J Babylon 2019;16:119-22.
 19. Al-Hilali KA. Prevalence of hypovitaminosis D in adult Iraqi people including postmenopausal women. Sci Res J 2016;4:53.
 20. Hasani NJ. Assessment of Vitamin D Level in Sera of Primary School Students in Babylon Province. Journal of University of Babylon for Pure and Applied Sciences 2019;27(2): 189-194.
 21. [Yaqoo HA](#), [Haddad NZ](#), [Jawad AM](#). Serum vitamin D level, measured by two methods, in a sample of normal subjects in Basrah : [The Medical Journal of Basrah University](#);2019,37(2):106-114.
 22. Garcia-Bailo B, Badawi A, El-Sohemy A. Vitamin D status and biomarkers of type 2 diabetes risk in an ethnically diverse population. Can J Diabetes 2011;35:432-3 [[Google Scholar](#)]
 23. Brenner DR, Arora P, Garcia-Bailo B, et al. The relationship between metabolic syndrome components and inflammatory markers among non-diabetic Canadian adults. J Diabetes Metab 2012.
 24. Da Costa LA, Arora P, Garcia-Bailo B. The association between obesity, cardiometabolic disease markers and innate immunity-related inflammation in Canadian adults. Diabetes Metab Syndrome Obesity 2012.
 25. Brenner DR, Arora P, Garcia-Bailo B, et al. Plasma vitamin D and risk of the metabolic syndrome in Canadians. Clin Invest Med 2011;34:E377-E384 [P
 26. Badawi A, Klip A, Haddad P, et al. Type 2 diabetes mellitus and inflammation: Prospects for biomarkers of risk and nutritional intervention. Diabetes Metab Syndr Obes 2010;26:3:173-86
 27. Badawi A, Arora P, Sadoun E, Al-Thani AA, Thani MH. Prevalence of vitamin d insufficiency in qatar: a systematic review. J Public Health Res. 2012 Dec 28;1(3):229-35.
 28. Forrest KY, Stuhldreher WL. Prevalence and correlates of vitamin D deficiency in US adults. Nutri Res (N Y, NY). 2011;31(1):48-54.
 29. Freishtat RJ, Iqbal SF, Pillai DK, Klein CJ, Ryan LM, Benton AS, Teach SJ. High prevalence of vitamin D deficiency among inner-city African American youth with asthma in Washington, DC. J Pediatr. 2010;156(6):948-52.
 30. Lim JU, Lee JH, Kim JS, Hwang YI, Kim TH, Lim SY, et al. Comparison of World Health Organization and Asia-Pacific body mass index classifications in COPD patients. Int J Chron Obstruct Pulmon Dis. 2017; 12, 2465-2475.
 31. Looker AC, Johnson CL, Lacher DA, et al. Vitamin D status: United States, 2001-2006. National Center for Health Statistics, Centers for Disease Control and Prevention. Data Brief Report n. 59, 2011.
 32. Al Anouti F, Thomas J, Abdel-Wareth L. Vitamin D deficiency and sun avoidance among university students at Abu Dhabi, United Arab Emirates. Dermatoendocrinol 2011;3:235-9
 33. Mithal A, Wahl DA, Bonjour JP. IOF Committee of Scientific Advisors (CSA) Nutrition Working Group 2009. Global vitamin D status and determinants of hypovitaminosis D. Osteoporos Int 2009;20:1807-20.

34. Fields J, Trivedi NJ, Horton E. Vitamin D in the Persian Gulf: Integrative Physiology and Socioeconomic Factors. *Curr Osteoporos Rep* 2011;9:243-50.
35. MUSAIGER OA, AL-HAZZAA HM, TAKRURI HR. Change in nutrition and lifestyle in the Eastern Mediterranean Region: Health impact. *J Nutr Metab* 2012. doi:10.1155/2012/43676.