

# Recent Advances in HIV Diagnosis (Orasure and Oraquick)

**B. Amrithaa<sup>1</sup>, R.V. Geetha<sup>2</sup>, Lakshmi.T<sup>3</sup>**

<sup>1</sup>Research Associate, <sup>2</sup>Associate Professor, Department of Microbiology, Saveetha Dental College, Saveetha Institute of Medical & Technical Science, Saveetha University, Chennai, India, <sup>3</sup>Associate Professor, Department of Pharmacology, saveetha Dental College, saveetha university, Chennai, India

## Abstract

Acquired Immunodeficiency syndrome(AIDS) is an infection caused by the Human Immunodeficiency Virus(HIV). The infection straight forwardly influences the cells of the insusceptible framework, especially the T lymphocytes, bringing about various sharp contaminations prompting the terminal condition AIDS. Since no antibodies are accessible for this shocking illness, HIV disease is a danger and can without much of a stretch be named as a revile upon mankind. The Centers for Disease Control and Prevention (CDC) and numerous general wellbeing boards of trustees advocate for the fast HIV test that is accessible. Tests that can distinguish HIV contamination biomarkers inside 30days of disease, when beginning safe reactions are mounted. The OraQuick ADVANCE Rapid HIV - 1/2 Antibody test is sensitive– utilizing subjective immunoassay to distinguish the antibodies of HIV-1 and HIV-2 out of an oral liquid. Orasure Technologies is enabling the worldwide network to improve wellbeing and health by giving access to precise fundamental data. OraQuick and Orasure determination are rapid, flexible, accurate, simple and patient preferred to perceive antibodies to all HIV-1 subtypes, especially in making nations.

**Keywords :** HIV, AIDS, Orasure, OraQuick, Rapid diagnostic kits

## Introduction

Acquired immunodeficiency syndrome(AIDS) is an infection caused by the human immunodeficiency infection(HIV). HIV contamination is an amazingly present danger and can without a very remarkable stretch be named as a berate upon mankind. Set of scientists initially observed and saw the closeness of AIDS as a real disease following an extension in the event of remarkable enterprising contamination and malignancies among regardless among in any case among otherwise healthy homosexual men<sup>1</sup>. The Centers for Disease Control and Prevention (CDC) and numerous general wellbeing council keep on pushing for the most delicate quick HIV

test that is accessible, both to recognize positive patients when they are at their generally irresistible, and to guarantee that negative patients are really negative when beginning a pre-exposure prophylaxis routine<sup>2</sup>. Tests that can identify HIV disease biomarkers within 30 days of disease, when initial immune responses are mounted<sup>3</sup>. The infected patient's IgM reaction is regularly used to identify intense disease inside a 20–25 days of window period after contamination <sup>4</sup>.

Quick HIV immunizer tests have altered HIV analysis by encouraging the testing of a large number of individuals around the world<sup>5</sup>. The accessibility of reasonable, purpose of administration HIV testing is particularly significant in low-pay, high HIV trouble nations which do not have the money related and innovative assets to perform progressively refined research center based assays<sup>6</sup>. OraSure Technologies is enabling the worldwide network to improve wellbeing and health by giving access to precise fundamental data. OraSure is a pioneer in the turn of events, assembling and dispersion of purpose of care demonstrative tests, sub-atomic assortment devices and different innovations intended to recognize or analyze basic ailments. Its first-

---

### Corresponding author:

#### **R.V. Geetha**

Associate Professor, Department of Microbiology  
Saveetha Dental College, Saveetha Institute of Medical  
and Technical Science, Saveetha University  
162, Poonamallee High Road Chennai-600077  
Tamil Nadu, India. Email : geetha@saveetha.com  
Phone number: 9710456203

to-showcase, imaginative items incorporate fast tests for the recognition of antibodies to HIV<sup>7</sup>.

### HIV Transmission

HIV might be transmitted through certain body liquids that are fit for containing high concentrations of HIV. These liquids incorporate blood, semen, vaginal and rectal emissions, and breast milk. Notwithstanding the various courses by which HIV-1 can be transmitted beginning with one individual then onto the following, there is usually a sorted out and reproducible appearance in the blood of viral and host markers of sickness following a clinically productive transmission event<sup>8,9</sup>. Transmit HIV through various specific activities<sup>10</sup>. Most regularly, individuals get or transmit HIV through sexual practices and needle or syringe use<sup>11</sup>. Just certain body liquids, blood serum, rectal liquids and breast milk from an individual who HIV can transmit<sup>12</sup>. These liquids must interact with mucous layer or harmed tissue or straightforwardly infused into the circulatory system.

### Epidemiology of HIV

The AIDS case announcing arrangement of the Centers of Disease Control and Prevention (CDC) is the main complete national population based information accessible to screen the epidemic<sup>13</sup>. Information is valuable in assessing ailment predominance and frequency, revealed AIDS cases are just clinical tips of something larger impacts created by HIV contaminations<sup>14</sup>. HIV seroprevalence reviews are data for their description of the size of the epidemic, however they speak to individuals whose date of contamination is unknown<sup>15</sup>. HIV rate information is for more data for observing the current course of the epidemic<sup>16</sup>. Since HIV contamination isn't reportable in all states and on the grounds that most investigations of HIV have excluded agent tests<sup>17</sup>. HIV observation gives data that is of restricted incentive in forecasting the fate of the epidemic<sup>18</sup>. Better developed social epidemiologic information known risk practices could give information about possible transmission and future spread<sup>19,20</sup>.

### Immune markers of HIV

When HIV has entered the body, the insusceptible framework starts hostile to HIV counter acting agent and binds to dendritic cells which convey the infection

to CD4+ T cells in lymphoid tissue establishing the disease<sup>21,22</sup>. Virus replication quickens delivering massive viremia and wide dispersal of virus all through the body's lymphoid tissues<sup>23</sup>. An immune reaction against the infection causes some protection yet a constant relentless contamination is set up. The creation of cytokines and cell divisions that direct the resistant reaction for protection likewise cause HIV Virus replication quickens conveying massive viremia and wide dispersal of infection all through the body's lymphoid tissues<sup>24</sup>. There is a rapid turnover of CD4+ T cells that ultimately leads to their destruction and to a change in lymphoid tissues that prevent immune reactions. Anyway it can take one to a half year for an individual exposed to HIV to produce measurable quantities of antibody<sup>25</sup>.

The immune reaction is weakened as memory t cells (CD4+CCR5+) are Destroyed. During development, HIV proteases divide the poly-proteins<sup>26,27</sup>. Into individual functional HIV proteins<sup>28</sup>. The mature various are able to cause another host cell. Maturation happens either in the forming buds or subsequent to maturing from the host cell. Tests to distinguish HIV with the nearness of antibodies to HIV, not simply the infection. Although a few tests are sensitive, there is a "window period". This is the period between the beginning of disease with HIV and the presence of distinguishable antibodies to the infection. On account of the most sensitive anti- HIV tests as of now suggested, the window period is around three weeks. This period might be longer if less sensitive tests are utilized<sup>29</sup>.

Recent advances in technology have led to various simple rapid tests being developed. Most of these tests come in a kit and require no reagent, equipment, training, or specified temperature controls, and tests can be performed at any time<sup>30</sup>. OraQuick and Orasure diagnosis are rapid, flexible, accurate, simple and patient preferred to recognize antibodies to all HIV-1 subtypes, particularly in creating countries.

### **ORAQUICK AND ORASURE**

The OraQuick® ADVANCE Rapid HIV-1/2 Antibody Test is a solitary use, subjective immunoassay to recognize antibodies of HIV-1 and HIV-2 in oral fluid, fingerstick whole blood, venipuncture whole blood and plasma specimens<sup>31,32</sup>. The OraQuick® ADVANCE

Rapid HIV-1/2 Antibody Test is normal for use as a condition-of-care test to help in the finish of tainting with HIV-1 and HIV-2. This test is proper for use in multi-test estimations proposed for quantifiable endorsement of quick HIV test results<sup>33</sup>. Clinical assessment of the OraQuick ADVANCE Rapid HIV-1/2 immunizer test has uncovered high affectability and explicitness. The test has numerous significant applications, expanding the open doors for willful directing and testing, and as a device for the scale-up of antiretroviral treatment in asset constrained settings<sup>34</sup>. Rapid, Affordable, quick, minimal effort, delicate, and explicit tests are expected to recognize antibodies to all HIV-1 subtypes, particularly in creating nations<sup>35</sup>. In addition in provincial regions assets are restricted, research center technicians much of the time inaccessible, phlebotomy is troublesome and socially less favored HIV test results can end up being extremely testing. Although oral liquid – based quick testing has a few alluring highlights and give helpful and feasible choice to blood – based testing<sup>36</sup>.

### **History of Orasure and Oraquick**

OraQuick is a HIV test made by Orasure Technologies and affirmed by the United States Food and Drug Administration in 2012. Orasure Technologies was chosen as one of the region's top places to work<sup>37</sup>. Perceived as a 2016 Top workplace, Orasure was only one of 15 organizations who were chosen among employers<sup>38</sup>. In 2011, Orasure gained DNA Genotek, a main supplier of oral liquid example collection, adjustment and planning products for molecular analytic applications. Furthermore, in 2012, Orasure got FDA endorsement for the Oraquick in-home HIV test, the first and only oral liquid fast OTC HIV test affirmed in the US and in 2019, Orasure gains CoreBiome, a microbiome administrations supplier, a urine sample collection device developer<sup>39</sup>.

Orasure and OraQuick recent diagnosis, which is very much grown these days. The definite determination of HIV contamination at any age requires finding testing that confirms the presence of HIV<sup>40</sup>. HIV disease definitely in children aged less than 18 months, measures are required to recognize the infection or its parts i.e Virological test<sup>41,42</sup>. Maternal HIV immune response is moved to the infant latently during pregnancy and afterward decays with a half – life 28-30 days in non

breastfed infants<sup>43,44</sup>.

### **OraQuick ADVANCE Rapid HIV-1/2 Antibody Test**

The OraQuick ADVANCE Test is FDA endorsed to recognize antibodies to HIV-1 and HIV-2 in entire blood, plasma, or oral fluid samples. The assay strip in the device consists of a membrane of nitrocellulose stripped with control and test lines<sup>45</sup>. The test line is striped with both HIV-1 and HIV-2 synthetic peptides and adjusted streptavidin (a biotin-binding protein). HIV- specific antibodies are presented with the sample, and, if present, will interact with biotinylated HIV-1 and HIV-2 peptides and colloidal gold marked with Protein A, which are impregnated in various segments of the device<sup>46</sup>. This sample reagent mixture then flows up the nitrocellulose, with complexed gold particles binding to the test line segments in different “sandwich” setups. The presence of HIV-1 as well as HIV-2 antibodies in the tested specimen is shown by the presence of a reddish purple coloured band at the test line area of the device within 20–40 minutes<sup>47</sup>.

### **Conclusion**

The review briefly outlines that OraQuick ADVANCE and Orasure HIV -1/2 Antibody Test detects antibodies to HIV-1 and HIV-2 in 20 minutes and this diagnosis is rapid, flexible, accurate, simple and patient preferred. This rapid test showed good sensitivity in detecting antibodies against HIV from oral samples making diagnosis easy. These tests detect IgM antibodies of acute infection within windows period based on capacity, and detect seroconversion approximately 20-25 days after infection..

**Acknowledgement:** We thank Saveetha dental college for the support given to conduct the study

**Conflict of Interest :** NIL

**Ethical Clearance:** It is taken from “Saveetha Institute Human Ethical Committee” (Ethical Approval Number- SDC/SIHEC/2020/DIASDATA/0619-0320)

### **References**

1. Bhatti AB, Usman M, Kandi V. Current Scenario of HIV/AIDS, Treatment Options, and Major Challenges with Compliance to Antiretroviral

- Therapy. *Cureus*. 2016 Mar 1;8(3):e515.
2. M. BB, Centers for Disease Control and Prevention (U.S.), Michele OS, G. WL, Berry B, G. WB, et al. Laboratory testing for the diagnosis of HIV infection : updated recommendations [Internet]. 2014. Available from: <http://dx.doi.org/10.15620/cdc.23447>
  3. Lewis JM, Macpherson P, Adams ER, Ochodo E, Sands A, Taegtmeier M. Field accuracy of fourth-generation rapid diagnostic tests for acute HIV-1 [Internet]. Vol. 29, *AIDS*. 2015. p. 2465–71. Available from: <http://dx.doi.org/10.1097/qad.0000000000000855>
  4. Martin IB, Williams V, Ferguson D, Read S. Performance of and preference for oral rapid HIV testing in The Bahamas [Internet]. Vol. 11, *Journal of Infection and Public Health*. 2018. p. 126–9. Available from: <http://dx.doi.org/10.1016/j.jiph.2017.06.005>
  5. Ashwin KS, Muralidharan NP. Vancomycin-resistant enterococcus (VRE) vs Methicillin-resistant Staphylococcus Aureus (MRSA). *Indian J Med Microbiol*. 2015 Feb;33 Suppl:166–7.
  6. Zachary D, Mwenge L, Muyoyeta M, Shanaube K, Schaap A, Bond V, et al. Field comparison of OraQuick® ADVANCE Rapid HIV-1/2 antibody test and two blood-based rapid HIV antibody tests in Zambia [Internet]. Vol. 12, *BMC Infectious Diseases*. 2012. Available from: <http://dx.doi.org/10.1186/1471-2334-12-183>
  7. Sanders RW. The HIV-1 Envelope Glycoproteins: Folding, Function and Vaccin Design. Amsterdam University Press; 2003. 337 p.
  8. Shaw GM, Hunter E. HIV transmission. *Cold Spring Harb Perspect Med* [Internet]. 2012 Nov 1;2(11). Available from: <http://dx.doi.org/10.1101/cshperspect.a006965>
  9. Vaishali M, Geetha RV. Antibacterial activity of Orange peel oil on Streptococcus mutans and Enterococcus-An In-vitro study. *J Adv Pharm Technol Res*. 2018;11(2):513.
  10. Priyadharsini JV, Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. In silico analysis of virulence genes in an emerging dental pathogen *A. baumannii* and related species [Internet]. Vol. 94, *Archives of Oral Biology*. 2018. p. 93–8. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2018.07.001>
  11. Shahzan MS, Sohaib Shahzan M, Smiline Girija AS, Vijayashree Priyadharsini J. A computational study targeting the mutated L321F of ERG11 gene in *C. albicans*, associated with fluconazole resistance with bioactive compounds from *Acacia nilotica* [Internet]. Vol. 29, *Journal de Mycologie Médicale*. 2019. p. 303–9. Available from: <http://dx.doi.org/10.1016/j.mycmed.2019.100899>
  12. Girija As S, Priyadharsini J V. CLSI based antibiogram profile and the detection of MDR and XDR strains of isolated from urine samples. *Med J Islam Repub Iran*. 2019 Feb 8;33:3.
  13. Smiline ASG, Vijayashree JP, Paramasivam A. Molecular characterization of plasmid-encoded blaTEM, blaSHV and blaCTX-M among extended spectrum  $\beta$ -lactamases [ESBLs] producing *Acinetobacter baumannii* [Internet]. Vol. 75, *British Journal of Biomedical Science*. 2018. p. 200–2. Available from: <http://dx.doi.org/10.1080/09674845.2018.1492207>
  14. Normand J, Vlahov D, Moses LE. Preventing HIV Transmission: The Role of Sterile Needles and Bleach. *National Academies*; 1995. 334 p.
  15. National Institute on Drug Abuse. Longitudinal Studies of HIV Infection in Intravenous Drug Users. 1991. 153 p.
  16. Girija ASS, Smiline Girija AS, Vijayashree Priyadharsini J, Paramasivam A. Plasmid-encoded resistance to trimethoprim/sulfamethoxazole mediated by dfrA1, dfrA5, sul1 and sul2 among *Acinetobacter baumannii* isolated from urine samples of patients with severe urinary tract infection [Internet]. Vol. 17, *Journal of Global Antimicrobial Resistance*. 2019. p. 145–6. Available from: <http://dx.doi.org/10.1016/j.jgar.2019.04.001>
  17. Girija SAS, Jayaseelan VP, Arumugam P. Prevalence of VIM- and GIM-producing *Acinetobacter baumannii* from patients with severe urinary tract infection [Internet]. Vol. 65, *Acta Microbiologica et Immunologica Hungarica*. 2018. p. 539–50. Available from: <http://dx.doi.org/10.1556/030.65.2018.038>
  18. Boulle A, Johnson L. Human immunodeficiency

- virus (HIV) infection [Internet]. *Infectious Disease Epidemiology*. 2016. p. 343–50. Available from: <http://dx.doi.org/10.1093/med/9780198719830.003.0025>
19. Inciardi JA. AIDS—A Strange Disease of Uncertain Origins [Internet]. Vol. 33, *American Behavioral Scientist*. 1990. p. 397–407. Available from: <http://dx.doi.org/10.1177/0002764290033004002>
  20. Maajida Aafreen M, Geetha RV, Lakshmi Thangavelu. Evaluation of anti-inflammatory action of *Laurus nobilis*-an in vitro study. *ijrps*. 2019 Apr 14;10(2):1209–13.
  21. Altfeld M, Fadda L, Frleta D, Bhardwaj N. DCs and NK cells: critical effectors in the immune response to HIV-1 [Internet]. Vol. 11, *Nature Reviews Immunology*. 2011. p. 176–86. Available from: <http://dx.doi.org/10.1038/nri2935>
  22. Priyadharsini JV, Vijayashree Priyadharsini J, Smiline Girija AS, Paramasivam A. An insight into the emergence of *Acinetobacter baumannii* as an oro-dental pathogen and its drug resistance gene profile – An in silico approach [Internet]. Vol. 4, *Heliyon*. 2018. p. e01051. Available from: <http://dx.doi.org/10.1016/j.heliyon.2018.e01051>
  23. Mahalingam S, Meanger J, Foster PS, Lidbury BA. The viral manipulation of the host cellular and immune environments to enhance propagation and survival: a focus on RNA viruses. *J Leukoc Biol*. 2002 Sep;72(3):429–39.
  24. Bieniasz PD. Intrinsic immunity: a front-line defense against viral attack [Internet]. Vol. 5, *Nature Immunology*. 2004. p. 1109–15. Available from: <http://dx.doi.org/10.1038/ni1125>
  25. Platt EJ, Durnin JP, Kabat D. Kinetic Factors Control Efficiencies of Cell Entry, Efficacies of Entry Inhibitors, and Mechanisms of Adaptation of Human Immunodeficiency Virus [Internet]. Vol. 79, *Journal of Virology*. 2005. p. 4347–56. Available from: <http://dx.doi.org/10.1128/jvi.79.7.4347-4356.2005>
  26. Paramasivam A, Vijayashree Priyadharsini J, Raghunandhakumar S. N6-adenosine methylation (m6A): a promising new molecular target in hypertension and cardiovascular diseases. *Hypertens Res*. 2020 Feb;43(2):153–4.
  27. Shahana RY, Muralidharan NP. Efficacy of mouth rinse in maintaining oral health of patients attending orthodontic clinics [Internet]. Vol. 9, *Research Journal of Pharmacy and Technology*. 2016. p. 1991. Available from: <http://dx.doi.org/10.5958/0974-360x.2016.00406.6>
  28. Scherdin U, Rhodes K, Breindl M. Transcriptionally active genome regions are preferred targets for retrovirus integration. *J Virol*. 1990 Feb;64(2):907–12.
  29. Greenwald JL, Burstein GR, Pincus J, Branson B. A rapid review of rapid HIV antibody tests. *Curr Infect Dis Rep*. 2006 Mar;8(2):125–31.
  30. Uttayamakul S, Likansakul S, Sunthornkachit R, Kuntiranont K, Louisirochanakul S, Chaovavanich A, et al. Usage of dried blood spots for molecular diagnosis and monitoring HIV-1 infection. *J Virol Methods*. 2005 Sep;128(1-2):128–34.
  31. Holguín Á, Gutiérrez M, Portocarrero N, Rivas P, Baquero M. Performance of OraQuick Advance® Rapid HIV-1/2 Antibody Test for detection of antibodies in oral fluid and serum/plasma in HIV-1 subjects carrying different HIV-1 subtypes and recombinant variants [Internet]. Vol. 45, *Journal of Clinical Virology*. 2009. p. 150–2. Available from: <http://dx.doi.org/10.1016/j.jcv.2009.04.002>
  32. Selvakumar R, Np M. COMPARISON IN BENEFITS OF HERBAL MOUTHWASHES WITH CHLORHEXIDINE MOUTHWASH: A REVIEW [Internet]. Vol. 10, *Asian Journal of Pharmaceutical and Clinical Research*. 2017. p. 3. Available from: <http://dx.doi.org/10.22159/ajpcr.2017.v10i2.13304>
  33. Martin M, Frese W, Lumsden C, Sandoval A. Building a Pediatric Oral Health Training Curriculum for Community Health Workers [Internet]. *Journal of Public Health Management and Practice*. 2017. p. 1. Available from: <http://dx.doi.org/10.1097/phh.0000000000000582>
  34. Alemnji GA, Ngulefac GA, Ndumbe PM, Asonganyi T. Field Evaluation of Calypte’s AWARETM Blood Serum Plasma (BSP) and Oral Mucosal Transudate (OMT) Rapid Tests for Detecting Antibodies to HIV-1 and 2 in Plasma

- and Oral Fluid [Internet]. Vol. 3, The Open AIDS Journal. 2009. p. 14–8. Available from: <http://dx.doi.org/10.2174/1874613600903010014>
35. Pant Pai N, Joshi R, Dogra S, Taksande B, Kalantri SP, Pai M, et al. Evaluation of diagnostic accuracy, feasibility and client preference for rapid oral fluid-based diagnosis of HIV infection in rural India. *PLoS One*. 2007 Apr 11;2(4):e367.
  36. Tucker JD, Bien CH, Peeling RW. Point-of-care testing for sexually transmitted infections: recent advances and implications for disease control. *Curr Opin Infect Dis*. 2013 Feb;26(1):73–9.
  37. Barakat M, University of Pennsylvania. HYDROGEOLOGICAL ASSESSMENT OF CAMEL'S HUMP FARM IN BETHLEHEM, PA [Internet]. 2018. Available from: <http://dx.doi.org/10.1130/abs/2018am-324790>
  38. Reynolds SJ, Ndongala LM, Luo C-C, Mwandagirwa K, Losoma AJ, Mwamba KJ, et al. Evaluation of a rapid test for the detection of antibodies to human immunodeficiency virus type 1 and 2 in the setting of multiple transmitted viral subtypes. *Int J STD AIDS*. 2002 Mar;13(3):171–3.
  39. Panel on Data and Research Priorities for Arresting AIDS in Sub-Saharan Africa. Preventing and Mitigating AIDS in Sub-Saharan Africa:: Research and Data Priorities for the Social and Behavioral Sciences. National Academy Press; 1996. 356 p.
  40. Parker M. HIV and Infant Feeding. (World Health Organization, Joint United Nations Programme on HIV/AIDS (UNAIDS) and United Nations Children's Fund, 1998.) US \$14.40 [Internet]. Vol. 32, *Journal of Biosocial Science*. 2000. p. 286–7. Available from: <http://dx.doi.org/10.1017/s0021932000222839>
  41. Nesheim S, Palumbo P, Sullivan K, Lee F, Vink P, Abrams E, et al. Quantitative RNA Testing for Diagnosis of HIV-Infected Infants [Internet]. Vol. 32, *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2003. p. 192–5. Available from: <http://dx.doi.org/10.1097/00126334-200302010-00011>
  42. Marickar RF, Geetha RV, Neelakantan P. Efficacy of Contemporary and Novel Intracanal Medicaments against *Enterococcus Faecalis* [Internet]. Vol. 39, *Journal of Clinical Pediatric Dentistry*. 2014. p. 47–50. Available from: <http://dx.doi.org/10.17796/jcpd.39.1.wmw9768314h56666>
  43. Institute NC, National Cancer Institute. World Health Organization Clinical Staging of HIV/AIDS for Adults and Adolescents Clinical Classification [Internet]. Definitions. 2020. Available from: <http://dx.doi.org/10.32388/ctil6a>
  44. Pratha AA, Ashwatha Pratha A, Geetha RV. Awareness on Hepatitis-B vaccination among dental students-A Questionnaire Survey [Internet]. Vol. 10, *Research Journal of Pharmacy and Technology*. 2017. p. 1360. Available from: <http://dx.doi.org/10.5958/0974-360x.2017.00240.2>
  45. Nkenfou CN, Kembou JT, Djikeng A, Domkam I, Tchuinkam T. An evaluation of human immunodeficiency virus oral screening test awareness and preferences in the West region of Cameroon [Internet]. Vol. 8, *Journal of Infection and Public Health*. 2015. p. 254–9. Available from: <http://dx.doi.org/10.1016/j.jiph.2014.10.004>
  46. Smallwood M, Vijn R, Nauche B, Lebouché B, Joseph L, Pant Pai N. Evaluation of a Rapid Point of Care Test for Detecting Acute and Established HIV Infection, and Examining the Role of Study Quality on Diagnostic Accuracy: A Bayesian Meta-Analysis. *PLoS One*. 2016 Feb 18;11(2):e0149592.
  47. Wesolowski LG, MacKellar DA, Facente SN, Dowling T, Ethridge SF, Zhu JH, et al. Post-marketing surveillance of OraQuick whole blood and oral fluid rapid HIV testing [Internet]. Vol. 20, *AIDS*. 2006. p. 1661–6. Available from: <http://dx.doi.org/10.1097/01.aids.0000238413.13442.ed>