

Global Burden of Periodontal Diseases: A Literature Review

Subhangi Chandan¹, Abhaya Chandra Das², Sourav Panda²,
Rashmita Nayak³, Rinkee Mohanty³, Manoj Kumar²

¹Intern, Institute of Dental Sciences, ²Reader, ³Professor, Department of Periodontics and Oral Implantology, Institute of Dental Sciences, Siksha 'O' Anusandhan, (Deemed to be University), Bhubaneswar, Odisha, India

Abstract

Periodontal disease is prevalent in both advanced and growing nations and affects about 20-50% of the world population. Several hazard factors inclusive of smoking, heredity, diabetes, bad oral hygiene, age, pressure and medication which is related to periodontal illness. The devaluation inside occurrence and popularity of periodontitis can decrease its correlated fundamental illness and diminish their economic impact on health care systems. Periodontitis is one of the most important diseases which is chronic and leads to death and disability worldwide. The speediest increase in the hardship of the chronic disease is acquiring in the growing nations.

Keywords: Periodontal disease; Global burden; Risk Factors; Economic impact; Treatment policies

Introduction

Periodontitis is a continuous intemperate disease and characterised through loss of periodontal ligament and elimination of neighbouring bone.¹ There are different analytic explanations of periodontal disease and it may be acute or chronic. Acute periodontitis are of limited period, quick onset and associated with pain. Early diagnosis and prompt treatment is necessary. Due to the deposition of dental plaque, chronic periodontitis occurs. Disintegration of the supporting tissues of the teeth occurs in periodontitis. Deepening of periodontal pockets and attachment loss is the sign of periodontal disease. A huge prevalence of periodontitis in adults and elder persons makes it a mutual well-being burden.

Prevalence of periodontal diseases: Periodontitis is the most accepted oral condition among human

populations.² The frequency and popularity of periodontitis diverge due to preference, case-mix and sites and number of teeth investigated. The World Health Organisation (WHO) has managed overall oral health data bank by the help of the community periodontal index (CPI).³ Community periodontal index varies from 0 to 4.⁹ Community periodontal index score 0 indicates an absence of periodontitis; score 1 indicates bleeding observed during or after probing; score 2 indicates during probing supra or subgingival calculus seen; score 3 indicates hollow pathological pocket of 4 or 5 mm (gingival margin within the black band of the probe); score 4 indicates pathological periodontal pocket of 6mm or more (the black band of the probe is not visible).⁴ Among adolescents, widespread of calculus and bleeding problem is higher in developing nations than developed nations. The distribution of teenagers with calculus deposits varies from 35% to 70% in advancing nations and 4% to 34% in advanced nations. In advanced countries periodontal pockets of 4-5mm seen in a higher percentage of individuals. Correspondingly in grown-up, calculus deposits varies from 14-17% in advanced countries and 36-63% in advancing countries. Higher distribution of elder individuals (65-75 years) show pathological pockets of 6mm or more is seen in grown-up populations in both advanced and advancing nations.⁵

Corresponding Author:

Dr. Abhaya Chandra Das

Reader, Department of Periodontics and Oral Implantology, Institute of Dental Sciences, Siksha 'O' Anusandhan, (Deemed to be University), Bhubaneswar, Odisha, India
e-mail: drabhaya2011@gmail.com

Factors related to periodontal diseases: A few risk factors raise the possibility of periodontal diseases which commit toward the scientific number implication of periodontal disease. The essential risk factors for serious periodontal disease are the adoption of tobacco, stress, malnourishment, extreme alcohol drink, diabetes mellitus and other systemic diseases.

Modifiable Risk Factors:

- 1. Smoking:** The greater critical possibility causes for periodontal disease is smoking. The devaluation in periodontitis pervasiveness is associated with a decrease in the rate of smoking. In smokers, there might be increased in the loss of alveolar bone along with tooth loss than the people having no smoking habit and smokers have low results of patterns of periodontitis.⁶
- 2. Bad oral hygiene:** Bad oral hygiene is connected with periodontitis, the need of legitimate brushing of teeth and different proportion of oral hygiene can support affidavit of bacteria and develop deposition of plaque on teeth and gums which can address an aspect for fiery changes in tissues which is present in the periodontal region.⁷ There is an articulated relationship between bad oral hygiene and expanded aggregation of plaque, huge commonness and expanded seriousness of periodontitis.⁸
- 3. Hormones:** Harmonization in females improve probability of periodontitis.⁸ Ladies may encounter aggravation of gingiva previously period and at the time of ovulation because of a significant matched in progesterone which hinders fix of collagen strands and creates dilation of blood vessels. Pregnant ladies as often as possible display gingival changes, gum disease, now and again limited development of gingival tissues.⁹ Luckily, these provocative changes vanish inside barely after conveyance beyond motivating tenacious harm to tissues present in periodontal region. Inadequacy of oestrogen diminishes the thickness of bone after the midlife crisis that can come full circle in alveoli misfortune or the long run collapsing of teeth.¹⁰
- 4. Diabetes:** People suffering from diabetes are more likely to develop periodontitis which can raise blood sugar level and various diabetic complications. Diabetes is the foundational ingredients for periodontitis which cause a significant job in commencement and movement of the disease.¹¹ Diabetes is related to tendon demolition which

therefore prompt loss of tooth. Gingival crevicular fluid and salivation have higher convergence of fiery go-between including extraordinary sorts of cytokines encompassed by diabetic patients with periodontal disease when contrasted with people not suffering from diabetes with periodontitis. Periodontitis is generally recognised a complexity of diabetes. People who don't control their diabetes, are especially at risk. Periodontitis can be a result of low diabetes management and associated with cardiovascular disease and strokes.

- 5. Drugs:** Helplessness to contaminations and periodontitis increases lessened salivary stream because of medication powerlessness to contamination.¹² Periodontal sickness strengthens when there is reduced salivary stream because of certain limit the progression of salivation and produce Xerostomia, incorporates anti-depressants, anti-cholinergic, anti-histamines and beta receptor antagonists.³⁰ Some medications like nifedipine, cyclosporine and phenytoin can initiate the irregular development of gingival tissues which much of the time confounds the fitting expulsion of plaque beneath amplified gingival mass and in this manner bother current periodontitis.
- 6. Weight:** It is obvious form proof that the pressure diminishes progression of slavery discharges that thusly upgrade formation of plaque. An additive relationship among weight scores and slavery pressure markers, calamity of tooth, attachment loss and pocket depth of 5-8mm. An investigation of 300 observational pieces shown that the rest of pressure is identified with a safe framework and diverse immunological changes happen in light of various upsetting events. The discouraged people have appeared to a greater focus of cortisol in gingival crevicular liquid and react inadequately to treat the periodontal structures. Scholastic pressure additionally brings about bad oral hygiene and gingival irritation with expanded grouping of interleukin-1beta.¹³

Static Risk Factors:

- 1. Maturity:** Periodontal treatment increments with processing maturity which is the reason for the huge predominance of periodontitis which examines among overpopulation. Maturity is related to periodontitis and attachment loss endure essentially huge among population matured 60 to

69yrs contrasted and gathering of grown-up 40-50yers.

- 2. Heredity:** Heredity is one of the most important components which is related to periodontal disease which causes a few people further defenceless to infection than others. The mind-boggling interactions of hereditary variables among ecological or segment factors is speculated to exhibit broad varieties encompassed by various genetic and indigenous populations.

Relationship of periodontitis with other systemic conditions:

1. Ischemic heart disease: Reliable group of proof clarifies the connection between ischemic heart disease and periodontitis. A systematic survey recognised that the periodontal infection abidehazard element for congestive heart failure and affiliation autonomous of more hazard elements for example smoking, diabetes mellitus and financial condition. An examination of imminent and review considers, it has discovered that periodontitis probably going to purpose a 19% expansion in danger of coronary heart disease and expansion in relation hazard ranges directed toward 44% encompassed by people matures 65 years and over. An older survey and investigation of examinations discovered that periodontal malady with expanded degrees of foundational markers of bacteria was related toischemic heart disease. Periodontal ailment affiliation with stroke and fringe corridors infection is significantly more grounded than coronary illness. Correspondingly, an examination of 29 investigations revealed merged chances proportions of 2.35 and polled corresponding danger of 1.34 that recommended people with periodontitis has more noteworthy chance and higher chances of creating coronary illness than those without periodontal disease.¹⁴

- 2. Metabolicailment:** There are bidirectional communication and teamwork among diabetes mellitus and periodontitis. An imminent associate examinations of 628 subjects along check out recognised that insulin-independent diabetes mellitus people among serious periodontitis had 3.2 occasions danger of fatality due to coronary illness contrasted with the people with no or on the other hand gentle periodontal sickness. An examination presumed that the periodontitis developed

glycemic control for any event for non-insulin diabetesmellitus. Deliberate survey gave proof to help the job of periodontitis in improvement of adult-onset diabetes and also it's difficulties. Logical writing reliably underpins a bidirectional communication among periodontal disease and insulin obstruction. It has contended that periodontitis fuels insulin oppositions, constantquality embroiled in pathophysiology of catabolic sickness and adult-onset diabetes mellitus. It has additionally proposed that periodontal medication can lessen insulin opposition in diabetes. A few orderly audits have proposed a relationship between heftiness and periodontal ailment and recognised hazardfactors for improvement of periodontal diseases.Heftiness is appeared directed toward increment oxidated worry in periodontal tissues what's more motivated towards destructions. Predominance of heftiness has expanded drastically among globe5land affiliation with periodontal disease requires consideration among human service suppliers to forestall the general medical problems.¹⁵⁻¹⁷

- 3. Unfavourable pregnancy result:** Periodontitis is identified with unfavourable pregnancy results which incorporate maternal disease,low birth weight, preterm birth, preeclampsia, mist biological and immunological factors are embroiled in hidden strictures. Below economic condition,smoking and other diseases are now known to be related with untimely birth; in any case,more as of late it was discovered that periodontal malady is likewise unequivocally connected with untimely birth episodes.¹⁸
- 4. Rheumatoid joint pain (Rheumatoid Arthritis):** Periodontal malady is common along rheumatoid joint pain patients and malady remains start immune system reaction in rheumatoid joint pain patients. It is recommended that periodontal malady and rheumatoid arthritis accept comparable hidden pathophysiology of disease. People suffering from rheumatoid arthritis have huge pervasiveness of bone obliteration and misfortune of toothwhichever additionally sequelae of periodontitis.
- 5. Respiratory illness:** The significance of keeping upideal oral consideration along individuals with incessant intrusive pneumonic ailment has underlined because of bidirectional relationship with periodontitis. Respiratory illness mainly includes asthma,pneumonia and lung cancer.

6. **Chronic Kidney Diseases:** There is a fractional connection between periodontitis and chronic kidney disease. Periodontal disease is a hazard factor for chronic kidney disease in an ideological investigation of 11,955 grown-ups precise audit of experimental and international examines found that patients along periodontal disease are expanded danger of chronic kidney disease and periodontitis brings about constructive appear in people with chronic kidney disease.¹⁹⁻²¹
7. **Diseases:** Hazard of oral cavity malignancy increments 5.23 occasion severe millimetre bone loss. Fitzpatrick saw that connection among periodontitis and vocal, abdominal, oesophageal, anorectic malignant growths have been accounted for all the more reliably in writing than with long and prostate malignant growths.
8. **Impedance of intellectual capacities:** More seasoned grown-up faces a decrease in their intellectual capacities which influence their practices including oral cleanliness habits. There is humble proof about a relationship between periodontal illness furthermore poor intellectual capacities as periodontal irritation has been appeared to influence discernment in old populations. Further, a late examination found that analytic attachment loss can advance beta gathering in mind which create intellectual dysfunction.

Determent of periodontitis:

1. **Avoidance of periodontal disease:** WHO suggests utilising incorporated general well-being preventive techniques which ought to be founded on regular hazard factor approach. Hazard factors, for example, stress, smoking, below economic conditions are real etc with periodontitis along with alternative fundamental interminable sickness, in this manner consideration of oral illness anticipation procedures in interminable fundamental ailment preventive activities can diminish the weight of malady in the vicinity of aligned of population.²²
2. **Cleanliness of mouthrehearses:** Appropriate cleaning of mouth, normal brushing of tooth and flossing are best among forestalling infection in mouth and periodontal disease. Despite most extreme importance of brushing of the tooth about the portion of populace brush two times a day. There are different diameter, configuration and sorts of brushes; be that as it may, two increasingly basic

sorts incorporated controlled toothbrushes and manual toothbrushes. Controlled toothbrushes often more bite of leeway up standard toothbrushes in lessening plaque present in the tooth.²³

3. **Starve:** Despite fact that the job of starving in anticipation of caries is increasingly huge contrasted and forestalling periodontal illness has been talked about in the writing. An eating regimen high in organic products, below in hefty and sugar is necessary for solid periodontal structures. Less calorie consumption is appeared directed toward lessening incendiary diversity and reduce the tissue harm in periodontitis.
4. **Fluoride utilisation:** Stannous fluoride has anti-plaque along with anti-gingivitis impacts what's more or lessen the extent of microscopic organisms and spirochetes in sub gingival zones, subsequently can assist with advancing gingival health.
5. **Utilization of antimicrobial specialist:** Chlorhexidine, triclosan, fundamental oils and mouthwashes and gels are utilised to prevent explicit microscopic organisms like plaque. Cholorhexidine diminishes dental plaque and gingival aggravation by bringing down provocation mediators. Gunsolley looked at the viability of anti-plaque and anti-gingivitis mouth rinses with oral cleanliness directions what's more grown-up prophylaxis and discovered extraordinary improvement in oral cleanliness.
6. **Smoking suspension:** Since smoking is a significant hazard factors for periodontal illness, in this way smoking end can forestall an extensive extend of periodontal cases. Smoking suspension not just further movement of periodontal ailment however can likewise decrease the periodontal tissue destruction. In light relationship among periodontitis and coronary heart infection, scaling has as of late appeared to lessen the frequency of intense cardiovascular infection localised necrosis also stroke.²⁴⁻²⁶

Conclusions

Whereas periodontitis is the common pervasive irresistible oral situation however it is stoppable and curable. Decrease in rate and pervasiveness of periodontal malady can bring about bringing down its related foundational ailments and confusions. Decreased periodontal ailment weight can limit treatment needs

and can lessen money related effect on wellbeing care frameworks. High commonness of periodontal ailment additionally requires the foundation of reconnaissance framework for oral ailments in the network. Preventive projects for periodontal ailment ought to use normal hazard way to deal with decrease the size of other incessant infections. Cost successful techniques would likewise improve interdisciplinary coordinated efforts among human service suppliers. Health care suppliers ought to be acquainted with Perio-fundamental connection and ought to have the option to analyse and allude patients to specific dental considerations to increase personal satisfaction of individuals. Research is expected to investigate fundamental instruments along with hazard variables of periodontitis and create imaginative preventive methodologies.

Ethical Permission: Not required

Conflict of Interests: None

Funding: None

References

- de Pablo P, Chapple IL, Buckley CD, Dietrich T. Periodontitis insystemic rheumatic diseases. *Nat Rev Rheumatol*2009; 5:218-24.
- Benjamin RM. Oral health: The silent epidemic. *Public Health Rep*2010;125:158-9.
- Ashby MT, Kreth J, Soundarajan M, Sivuilu LS. Influence of a modelhuman defensive peroxidase system on oral streptococcal antagonism. *Microbiology* 2009; 155:3691-700.
- Raitapuro-Murray T, Molleson TI, Hughes FJ. The prevalence ofperiodontal disease in a Romano-British population c 200-400 AD. *BrDent J* 2014; 217:459-66.
- Locker D, Slade GD, Murray H. Epidemiology of periodontal disease among older adults: A review. *Periodontol* 2000 1998; 16:16-33.
- Health Canada. Report on the Findings of the Oral Health Componentof the Canadian Health Measures Survey; 2007-2009.
- Eke PI, Dye BA, Wei L, et al. CDCPeriodontal Disease Surveillance workgroup: James Beck, et al. *J DentRes* 2012; 91:914-20.
- World Health Organization. WHO Global Oral Health Data; 2005.
- Sanz M, D’Aiuto F, Deanfield J, Fernandez-Avilés F. Europeanworkshop in periodontal health and cardiovascular disease-scientificevidence on the association between periodontal and cardiovasculardiseases: A review of the literature. *Eur Heart J Suppl* 2010;12 Suppl B: B3-12.
- Bergstrom J. Smoking rate and periodontal disease prevalence: 40-yeartrends in Sweden 1970-2010. *J ClinPeriodontol*2014; 41:952-7.
- Underner M, Maes I, Urban T, Meurice JC. Effects of smoking onperiodontal disease. *RevMal Respir*2009; 26:1057-73.
- Johnson GK, Hill M. Cigarette smoking and the periodontal patient.*J Periodontol* 2004; 75:196-209.
- Al Bandar JM, et al. Cigar, pipe, andcigarette smoking as risk factors for periodontal disease and tooth loss.*J Periodontol*2000; 71:1874-81.
- Reners M, Brex M. Stress and periodontal disease. *Int J Dent Hyg* 2007; 5:199-204.
- de Oliveira C, Watt R, Hamer M. Toothbrushing, inflammation, andrisk of cardiovascular disease: Results from Scottish Health Survey.*BMJ* 2010;340:c2451.
- Albandar JM. Global risk factors and risk indicators for periodontaldiseases. *Periodontol* 2000 2002; 29:177-206.
- Axelsson P, Lindhe J, Nyström B. On the prevention of caries andperiodontal disease. Results of a 15-year longitudinal study in adults.*J ClinPeriodontol*1991; 18:182-9.
- Güncü GN, Tözüm TF, Caglayan F. Effects of endogenous sexhormones on the periodontium - Review of literature. *Aust Dent J* 2005; 50:138-45.
- Markou E, Boura E, Tsalikis L, et al. Theinfluence of sex hormones on proinflammatory cytokines in gingivaof periodontally healthy premenopausal women. *J Periodontal Res* 2011; 46:528-32.
- Laine MA. Effect of pregnancy on periodontal and dental health. *ActaOdontolScand*2002; 60:257-64.
- Casanova L, Hughes FJ, Preshaw PM. Diabetes and periodontaldisease: A two-way relationship. *Br Dent J* 2014; 217:433-7.
- Chávarry NG, Vettore MV, Sansone C, Sheiham A. The relationshipbetween diabetes mellitus and destructive periodontal disease: A meta-analysis. *Oral Health Prev Dent* 2009; 7:107-27.
- Preshaw PM, Bissett SM. Periodontitis: An oral

- complication of diabetes. *Endocrinol Metab Clin North Am* 2013; 42:849-67.
24. Patel MH, Kumar JV, Moss ME. Diabetes and tooth loss: An analysis of data from the national health and nutrition examination survey, 2003-2004. *J Am Dent Assoc* 2013; 144:478-85.
25. Chapple IL, Genco R; Working group 2 of the joint EFP/AAP workshop. Diabetes and periodontal diseases: Consensus report of the Joint EFP/AAP Workshop on Periodontitis and Systemic Diseases. *J Periodontol* 2013; 84 4 Suppl: S106-12.
26. Scully C. Drug effects on salivary glands: Dry mouth. *Oral Dis* 2003; 9:165-76.