

Detrimental Effect of Alprazolam on Sperm Viability of Mice (*Mus musculus*) Exposed to Chronic Unpredictable Mild Stress

Intan Indah Permatasari¹, Reny I'tishom², Hanik Badriyah Hidayati³, Irwanto⁴

¹Researcher, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, ²Associate Professor, Department of Medical Biology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, ³Associate Professor, Department of Neurology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia, ⁴Professor, Department of Child Health, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

Abstract

Objective: This study aims to investigate the effect of alprazolam administration on sperm viability of mice (*Mus musculus*) strain Balb/C following CUMS for 53 days. **Method:** This study used a true experimental design through randomization. Control group kept undisturbed without treatment and therapy throughout 60 days. K1 and K2 were exposed to a 53 day period of CUMS along with 7-day pretreatment adaptation. K2 was adapted for a week, stressed for 53 days along with cotreatment of alprazolam (4 mg/kg BW). **Results:** K1 and K2 showed insignificant improvement of sperm viability ($p > 0.05$). Meanwhile, K0 and K1 showed significant value compared to K2 ($p < 0.05$). K2 showed little increase of sperm viability in CUMS group along with subsequent treatment of alprazolam. **Conclusion:** Overall, the results indicate an insignificant effect of alprazolam on sperm viability of CUMS-induced mice.

Keywords: Chronic Unpredictable Mild Stress, Alprazolam, Viability, Sperm

Introduction

Psychological stress like anxiety can be defined as an uncomfortable emotional experience due to the biochemical and psychological stress and behaviour. In dealing with stress, benzodiazepine drugs are often prescribed for hypnotic, sedative and anxiolytic symptoms at all ages in health services. As one of benzodiazepine group, alprazolam is ranked as 13th place among drugs that are commonly sold in 2012⁽¹⁾. Alprazolam is prescribed to treat panic disorder and anxiety. Further consumption in Indonesia during 2010-2013 increases with the average rate around 0.56 of Define Daily Doses for Statistical Purposes (S-DDD)⁽²⁾. One of the factors contributing into the increased consumption of drug is the pharmacological profile. On a pharmacodynamic property, alprazolam is more potent than that of diazepam with 1 mg of alprazolam relatively

equivalent to 10 mg of diazepam⁽⁴⁾. Additionally, it is well absorbed and metabolized extensively into the whole body after oral administration⁽⁵⁾. Thus, the pharmacological profile increases the prolonged drug use.

Few studies show that longer term use of alprazolam causes withdrawal syndrome, tolerance, and dependence⁽⁶⁾. Longer term use of alprazolam eventually disrupts the mechanism of chronic stress due to the activation of sympathetic adrenal system, hypothalamic-pituitary-gonadal (HPG) axis, and hypothalamic-pituitary-adrenal (HPA) axis. Furthermore, alprazolam can lead to side effects on the testes of mice. Studies showed that injection of alprazolam in mice possess a deleterious effect on structural histopathology in the testes and infertility. Infertility is evident on late spermatids with the head deformities⁽⁷⁾. Also, administration of alprazolam limits the intratesticular vascularization effecting on sperm evaluation⁽⁸⁾.

Corresponding author:

Reny I'tishom

Email: ritishom@fk.unair.ac.id

Chronic unpredictable mild stress (CUMS) has been used in the animal setting because it can potentially

stimulate depression⁽⁹⁾. Given the roles of HPA axis and HPG axis in the production of sperm under chronic stress conditions, the viable sperm quality in mice was evaluated in CUMS-induced mice in this present study.

Despite the scant research about alprazolam effect on male fertility, the chronic use of alprazolam still remains extensive. This present study was designed to investigate the effect of alprazolam administration on the sperm viability in CUMS-induced mice.

Materials and Method

Experimental Animals:

Thirty-three healthy male mice (*Mus musculus*) Balb/C, aged 6-8 week (25-30 gram) were used in this study and housed in animal laboratory, Faculty of Science and Technology, Universitas Airlangga under controlled conditions. Mice were acclimated for 7 days in prior to actual experiments. Each group consisting of 11 mice was obtained through randomization. Mice were maintained under a 12 hour light/dark cycle with 40-70% of relative humidity and housed per cage (40 cm length x 30 cm width x 18 cm height) with a close fitting wire, tap water and foods. The experiment protocols in this study have been approved by Ethical Eligibility and Research Committee (No. 296/EC/FKUA/2019, Faculty of Medicine, Universitas Airlangga).

Table1. Time length of stressors in CUMS protocol for 53 days^{(10),(11)}

Types of Stressors	Description	Time
Bath	The cage is not provided with mat base and the water is poured into the cage with normal temperature of ± 2 cm	30 minutes – 1 hour
Damp bedding	The mat is placed in the wet cage	2.5 hours – 3 hours
No bedding	The cage is not provided with mat base to expose the mice on the plastic as a base of the cage	3.5 hours
Bedding change	Bed cages is replaced with the reused bedding from other species object such as rat	Change per 15 – 30 minutes for 1.5 hours – 3 hours
Cage tilt	The cages with no base are tilted at an angle of 45 degrees in the hope that the mice gather in the cage corner.	1 hour – 2.5 hours
Altered light cycle	Exposure to a 5 watt light, then continuous alteration on and off.	Altered per 15 minutes during 1 hour – 3 hours
Social stress	Mice are transferred to the cages of other mice in order to expose the mice on the new environment and leave the mice alone without any social interaction	30 minutes – 60 minutes

Table 2. Type of interventions in CUMS protocol^{(10),(11)}

Monday	Tuesday	Wednesday	Thursday	Friday
Cagetilt (2.5 h)	No bedding (3.5 h)	Cagetilt (2.5 h)	Bathing (1 hour)	No bedding (2 h)
Dampbedding (2.5 h)	Socialstress (2 h)	Bathing (30 min)	Beddingchanges (every 30 minutesfor 3 hours)	Dampbedding (2 h)
Bathing (30 min)	Dampbedding (2,5 h)	Dampbedding (3 h)	Socialstress (30 min)	Cagetilt (1 h)
Socialstress (1.5 hour)	Light/dark (every 15 minutes carried out simultaneouslywithdampbedding)	Light/dark (1 hour and carried out simultaneously withdampbedding)	Light/dark (3 hours and carried out simultaneously withsocialstress)	Light/dark (2 hours and simultaneously carried out withBathing)
		Socialstress (1 hour)		Bathing (2 h)

CUMS Procedure:

The experimental group (K1, K2) were exposed to various stressors of CUMS model for 53 days according to the protocols described by Zhu et al. and de Andrade et al.,^{(11),(10)}with some minor modifications. The stress groups were subdivided into 4 groups (K1-1; K1-2; K2-1; K2-2). Each stress protocols wasdone for 7 hours (8 a.m. to 3 p.m.) at one time per day to prevent the mice from adapting.

Preparation of alprazolam:

Alprazolam was obtained as tablets in maximum dose (4 mg/kg BW). The applied dose of alprazolam was prepared by diluting the drugs in distilled water to 0.025 mg/kg/daily (equivalent to the therapeutic dose calculation for humans by Nair and Jacob)⁽¹²⁾. Alprazolam was suspended with 2% sodium CMC. Suspension of the mixture (0.1 ml) wasadministered to CUMS-induced mice through oral gavage. Control group was administered withdistilled water.

Viability Test:

To determine whether spermatozoa was alive or dead, observation was conducted by mixing one drop of

fresh semen with HE 0.5% on an object glass with cover glass.Each mean of alive or dead sperm with absorption of eosin was observed and compared.

Statistical Analysis

The data results were statistically assessed using one way ANOVA, Lavene’s test and Least Significant Difference (LSD) Post Hoc test. Statistical analysis was constructed using SPSS 25. Data distribution with p-value <0.05 was considered statistically normal.

Results

Based on the statistical findings, tests of alprazolam effect on CUMS-induced group for the period of CUMS (day 7 to day 21) showed no significant effect of alprazolam among K0, K1 and K2 groups. To assess data distribution, Saphiro Wilk test was used. It resulted in normal distribution with the value of p> 0.05. The data homogeneity and variance, Lavene’s test, One Way Anova, and LSD Post Hoc test were used. The data was homogeneous with p> 0.05. Out of 33 samples, the table showed there was a significant difference in between K0, K1, and K2 with p = 0.971 (p > 0.05) using One Way AnovaLavene’s test. The LSD Post Hoc test was then performed to determine significant differences among

groups. The statistical data analysis showed that K0 was 75.09% higher than K1 and K2 groups (Table 3). As a result, no significant difference was seen in the sperm viability data at 65.73% and 65.91%.

Table 3. Mean, Distribution, and variance of sperm viability data

Sample groups	Mean	SD	Minimum	Maximum	Normality (Sapphiro-Wilk)
K0 (control group)	75.09	5.873	65	84	0.772
K1 (CUMS-induced group)	65.73	6.498	54	74	0.325
K2 (CUMS + alprazolam treated group)	65.91	6.818	57	80	0.527

Table 4. Variance of sperm viability data using One Way Anova

	P value	Result
Control, CUMS-induced, CUMS + alprazolam treated groups	0.971	Significant

Table 5. Results of difference in sperm viability among groups (K0, K1, and K2)

	K1(CUMS-induced group)	K2(CUMS + alprazolam group)
K0 (control group)	0.002	0.002
K1(CUMS-induced group)	-	0.947

Discussion

Stress disrupts the hormonal regulatory system through HPG axis pathway as a major influence on infertility. Our current study showed that 53 days exposure to CUMS is sufficient to induce significant chronic stress and decrease the mean of sperm viability. Intermittent chronic stress, heat stress, obesity, and social defeat stress have been shown to have a suppressive effect on spermatogenesis⁽¹³⁻¹⁶⁾. However, after 42 days of alprazolam administration (5 mg/kg BW) and exposure to stressors, the chronic moderate stress-induced mice were observed. It has been found to elicit a neutralizing response against the damaging effects of moderate chronic stress, restore antioxidant balance and

reduce serum cortisol⁽¹⁷⁾.

This current study also correlates with the study using 8 different stressors for 6 weeks, showing an increase in cycle cell arrest of spermatogonia and apoptosis of spermatid cells. There was also an increase of glucocorticoid expression in spermatogonia and spermatocytes due to the feedback mechanism of the HPA and HPG axis pathways⁽¹⁸⁾. The germ cells in the testes are particularly susceptible to ROS because of the high levels of polyunsaturated fatty acids and low antioxidant capacity⁽¹⁹⁻²¹⁾.

It is evident that ROS levels result in the formation of abnormal sperm morphology. ROS as a signal has the

potential to induce apoptosis^(22,23). The capacitation, acrosome reactions and binding of spermatozoa to zona pellucida can take place at low levels of ROS. The processes would not occur when the production of ROS exceeds normal limits, resulting in infertility. The male infertility is manifested in decreased sperm quality such as motility, viability, and defects in spermatozoa-oocyte fusion^(24,25).

Sperm viability is associated with the intact plasma membrane due to its interaction between spermatozoa and oocyte⁽²⁶⁾. Aside from that, this current study is in line with the previous study that observed the decreased sperm quality in diazepam-treated mice with various doses (2 mg/kg BW; 5 mg/kg BW; 10 mg/kg BW) for 8 weeks⁽²⁷⁾. In diazepam-treated mice (5 mg/kg BW), the significant decreased sperm was heightened as a consequence of antioxidant imbalance and lower level of testosterone.

To assess the drug toxicity, the morphology of spermatozoa can be an important aspect of consideration. Low testosterone levels disturb the differentiation of spermatids after spermiogenesis⁽²⁸⁻³⁰⁾. Meanwhile, oral administration of fluoxetine in CUMS-induced mice resulted in a decrease of sperm concentration, sperm motility and an increased number of abnormal spermatozoa⁽³¹⁾. Administration of fluoxetine (200 mg/kg) in male rats for 60 days resulted in a significant decrease in spermatogenesis, levels of FSH, the weight of testes, epididymides, ventral prostate and seminal vesicle⁽³²⁾. Other clinical study proved the effect of paroxetine administration in 35 healthy male volunteers for 5 weeks. The results showed that there was an increase in abnormal DNA fragmentation considerably⁽³³⁾.

On the other hand, oral administration of alprazolam (0.5 mg/kg BW, 1.5 mg/kg BW, 4.5 mg/kg BW) in mice for 3 months caused gradual decrease in serum testosterone, significantly at a dose of 4.5 mg/kg BW⁽⁷⁾. Corresponding to the previous studies, the dosage (4 mg/kg body weight) in this study was relatively maximum in humans.

Nevertheless, another study found out that alprazolam exerts a therapeutic effect on a 6-hour trial of acute immobilization stress-induced mice⁽³⁴⁾. Because of

HPA axis pathway activation, it has an impact on several biological effects either at the central or peripheral levels. In this case, pretreatment with alprazolam (0.25 and 0.5 mg/kg/ BW) elicits a reversible effect of anxiety, analgesia, and impaired locomotor activity. It was effective to produce neuroprotective effects⁽³⁴⁾. Positive effects in chronic moderate stress-induced mice within 21 days were against the damaging outcomes of moderate chronic stress, antioxidant balance restoration, and serum cortisol reduction after treated with alprazolam (5 mg/kg/ BW)⁽¹⁷⁾. On the contrary, K2 group in this current study had a progressive build-up of oxidative stress due to CUMS.

In fact, the present study in K2 group at the same dose (4 mg / kg BW) as in human doses has not been shown to be able to suppress viable sperm levels by <50%. The standard cut-off value for human semen viability should be at least 58% of alive sperm⁽³⁵⁾. Semen that contains alive sperm below 50% is identified as abnormal when observed and assessed. The alprazolam administration at maximum dose does not reflect that it should be considered as toxic. It is interesting to point that a change of antioxidant capacity was also observed in rats exposed to immobilization, cold and water immersion for 21 days^(36,37). Exposure to chronic stress for 40 days which increases lipid peroxidation and protein oxidation in the cortex area may also explain why alprazolam does not improve the sperm quality^(38,39).

Conclusion

Our findings suggest that alprazolam administration decreases sperm viability of CUMS-induced mice (*Mus musculus*). Further studies are necessary to investigate about the effect of alprazolam with various doses on sperm quality of mice (*Mus musculus*) following CUMS.

Ethical Clearance: This study was confirmed by Ethical Eligibility and Ethical Research Committee, Faculty of Medicine, Universitas Airlangga.

Conflict of Interest: The authors report no conflict of interest.

Source of Funding: Self funding.

Acknowledgments: The authors wish to thank the

members at Department of Medical Biology, Faculty of Medicine and animal laboratory section, Faculty of Science and Technology, Universitas Airlangga for providing the laboratory and experimental facilities.

References

- Grohol JM. Top 25 psychiatric medication prescriptions for 2011 [Internet]. 2011. Available from: <https://psychcentral.com/lib/top-25-psychiatric-medication-prescriptions-for-2013#1>
- IMS Health. Top 25 medicines by dispensed prescriptions (US). 2013.
- Priyatni N. Berapa Kebutuhan Diazepam Untuk Memenuhi Pelayanan Kesehatan Di Indonesia? Studi Kasus Konsumsi Diazepam Di Indonesia. *J Manaj Pelayanan Inf*. 2016;6:297–302.
- National Collaborating Centre for Mental Health. Antenatal and postnatal mental health: The NICE Guideline on clinical management and service guidance—updated edition. *Br Psychol Soc R Coll Psychiatr*. 2014;
- Ait-Daoud N, Hamby A, Sharma S, Blevins D. A Review of Alprazolam Use, Misuse, and withdrawal. *J Addict Med*. 2017;12:4–10.
- Banerjee I, Roy B, Sathian B, Kumar S, Saha A. Medications for Anxiety: A Drug utilization study in Psychiatry Inpatients from a Tertiary Care Centre of Western Nepal. *Nepal J Epidemiol*. 2011;1:119–25.
- Attia, A. Histological and Ultrastructural Studies of The Effect of Alprazolam-Induced Testicular Damage in Male Mice. *Egypt Soc Exp Biol*. 2013;9:1–14.
- Semet M, Paci M, Saïas-Magnan J, Metzler-Guillemain C, Boissier R, Lejeune H, et al. The Impact of drugs on male fertility: a review. *Andrology*. 2017;5:640–63.
- López-López AL, Jaime HB, Escobar Villanueva MDC, Padilla MB, Palacios GV AF. Chronic unpredictable mild stress generates oxidative stress and systemic inflammation in rats. *Physiol Behav*. 2016;161:15–23.
- de Andrade JS, Céspedes IC, Abrão RO, da Silva JM, Ceneviva R, Ribeiro DA, et al. Effects of acute restraint and unpredictable chronic mild stress on brain corticotrophin releasing factor mRNA in the elevated T-maze. *Behav Brain Res*. 2018 Jan;337:139–50.
- Zhu S, Shi R, Wang J, Wang J-F, Li X-M. Unpredictable chronic mild stress not chronic restraint stress induces depressive behaviours in mice. *Neuroreport*. 2014 Oct;25(14):1151–5.
- Nair AB, Jacob S. A simple practice guide for dose conversion between animals and human. *J basic Clin Pharm*. 2016 Mar;7(2):27–31.
- Nirupama M, Devaki M, Nirupama R, Yajurvedi HN. Chronic intermittent stress-induced alterations in the spermatogenesis and antioxidant status of the testis are irreversible in albino rat. *J Physiol Biochem*. 2013 Mar;69(1):59–68.
- Durairajanayagam D, Agarwal A, Ong C. Causes, effects and molecular mechanisms of testicular heat stress. *Reprod Biomed Online*. 2015 Jan;30(1):14–27.
- Wang B, Zhou J, Zhuang Y-Y, Wang L-L, Pu J-X, Huang Y-H, et al. The Non-Peptide Vasopressin V1b Receptor Antagonist, SSR149415, Ameliorates Spermatogenesis Function in a Mouse Model of Chronic Social Defeat Stress. *J Cell Biochem*. 2017 Nov;118(11):3891–8.
- Jia Y-F, Feng Q, Ge Z-Y, Guo Y, Zhou F, Zhang K-S, et al. Obesity impairs male fertility through long-term effects on spermatogenesis. *BMC Urol*. 2018 May;18(1):42.
- Kori RS, Aladakatti RH, Desai SD, Das KK. Effect of drug alprazolam on restrained stress induced alteration of serum cortisol and antioxidant vitamins (Vitamin C and E) in male albino rats. *J Clin Diagnostic Res*. 2016;10(8):AF07–9.
- Zou P, Wang X, Yang W, Liu C, Chen Q, Yang H, et al. Mechanisms of stress-induced spermatogenesis impairment in male rats following unpredictable chronic mild stress (uCMS). *Int J Mol Sci*. 2019;20(18).
- Vernet P, Aitken RJ, Drevet JR. Antioxidant strategies in the epididymis. *Mol Cell Endocrinol* [Internet]. 2004;216(1):31–9. Available from: <http://www.sciencedirect.com/science/article/pii/S0303720703004532>
- Aitken RJ, Roman SD. Antioxidant Systems and Oxidative Stress in the Testes. *Oxid Med Cell Longev* [Internet]. 2008;1:616821. Available from: <https://doi.org/10.4161/oxim.1.1.6843>
- Agarwal A, Virk G, Ong C, du Plessis SS. Effect of oxidative stress on male reproduction. *World J*

- Mens Health. 2014 Apr;32(1):1–17.
22. Vaithinathan S, Saradha B, Mathur PP. Methoxychlor induces apoptosis via mitochondria- and FasL-mediated pathways in adult rat testis. *Chem Biol Interact* [Internet]. 2010;185(2):110–8. Available from: <http://www.sciencedirect.com/science/article/pii/S0009279710001535>
 23. Aly HAA, Khafagy RM. Taurine reverses endosulfan-induced oxidative stress and apoptosis in adult rat testis. *Food Chem Toxicol* [Internet]. 2014;64:1–9. Available from: <http://www.sciencedirect.com/science/article/pii/S0278691513007527>
 24. Sikka SC. Relative impact of oxidative stress on male reproductive function. *Curr Med Chem*. 2001 Jun;8(7):851–62.
 25. Sheweita SA, Tilmisany AM, Al-Sawaf H. Mechanisms of male infertility: role of antioxidants. *Curr Drug Metab*. 2005 Oct;6(5):495–501.
 26. Khatun A, Rahman MS, Pang MG. Clinical assessment of the male fertility. Vol. 61, *Obstetrics and Gynecology Science*. Korean Society of Obstetrics and Gynecology; 2018. p. 179–91.
 27. Taher, MA, Anber Z. Effect of Diazepam on The Male Reproductive System in Male Rats. *World J Pharm Pharm Sci*. 2015;14:60–78.
 28. Chauvigné F, Zapater C, Gasol JM, Cerdà J. Germ-line activation of the luteinizing hormone receptor directly drives spermiogenesis in a nonmammalian vertebrate. *Proc Natl Acad Sci* [Internet]. 2014 Jan 28;111(4):1427 LP – 1432. Available from: <http://www.pnas.org/content/111/4/1427.abstract>
 29. Ramaswamy S, Weinbauer GF. Endocrine control of spermatogenesis: Role of FSH and LH/ testosterone. *Spermatogenesis*. 2014 Mar 4;4(2):e996025.
 30. Martins FF, Aguila MB, Mandarim-de-Lacerda CA. Impaired steroidogenesis in the testis of leptin-deficient mice (ob/ob -/-). *Acta Histochem* [Internet]. 2017;119(5):508–15. Available from: <http://www.sciencedirect.com/science/article/pii/S0065128116303695>
 31. Sakr HF, Abbas AM, Elsamanoudy AZ, Ghoneim FM. Effect of fluoxetine and resveratrol on testicular functions and oxidative stress in a rat model of chronic mild stress-induced depression. *J Physiol Pharmacol an Off J Polish Physiol Soc*. 2015 Aug;66(4):515–27.
 32. Bataineh HN, Daradka T. Effects of long-term use of fluoxetine on fertility parameters in adults male rats. *Neuroendocrinol Lett* [Internet]. 2007 [cited 2020 Jun 17];28(3):321–5. Available from: <https://pubmed.ncbi.nlm.nih.gov/17627270/>
 33. Tanrikut C, Feldman AS, Altemus M, Paduch DA, Schlegel PN. Adverse effect of paroxetine on sperm. *Fertil Steril*. 2010 Aug;94(3):1021–6.
 34. Goyal R, Kumar A. Protective effect of alprazolam in acute immobilization stress-induced certain behavioral and biochemical alterations in mice. *Pharmacol Reports*. 2007;59(3):284–90.
 35. World Health Organization. Laboratory manual for the examination and processing of human semen [Internet]. Cambridge: Cambridge University Press. 2010 [cited 2020 Jun 19]. p. 32–99. Available from: http://whqlibdoc.who.int/publications/2010/9789241547789_eng.pdf
 36. Sahin E, Gümüşlü S. Immobilization stress in rat tissues: alterations in protein oxidation, lipid peroxidation and antioxidant defense system. *Comp Biochem Physiol C Toxicol Pharmacol*. 2007 Jan;144(4):342–7.
 37. Zafir A, Banu N. Induction of oxidative stress by restraint stress and corticosterone treatments in rats. *Indian J Biochem Biophys*. 2009 Feb;46(1):53–8.
 38. Lucca G, Comim CM, Valvassori SS, Réus GZ, Vuolo F, Petronilho F, et al. Effects of chronic mild stress on the oxidative parameters in the rat brain. *Neurochem Int*. 2009;54(5–6):358–62.
 39. López-López AL, Jaime HB, Escobar Villanueva MDC, Padilla MB, Palacios GV, Aguilar FJA. Chronic unpredictable mild stress generates oxidative stress and systemic inflammation in rats. *Physiol Behav*. 2016 Jul;161:15–23.