

Psychosocial Consequences of Children with Idiopathic Growth Hormone Deficiency in Baghdad

Adraa H. Shawq¹, Zaid W. Ajil¹

¹Lecturer, Pediatric Nursing Department, College of nursing, University of Baghdad, Iraq

Abstract

Study aim: to evaluate psychosocial problems of children with growth hormone deficiency.

Methodology: A cross sectional study, of (80) children from both gender, (4 – 15) years diagnosed with idiopathic growth hormone deficiency, attending endocrine clinics in Baghdad city. Consent form taken from children and their guardians to participate in the study. Pediatric Psychosocial Symptoms Inventory (mood, behavioural, somatic, learning, and personality problems) was used to evaluate children psychosocial consequences. SPSS programme version 23 used for data analyses.

Result: children with growth hormone deficiency experience moderate psychosocial problems, especially personality and behavioural problems.

Recommendation: The researchers recommended details psychological assessment for children with growth hormone deficiency during their routine physical examination and provide suitable psychological support for them and their parent.

Key words: *psychosocial consequences, growth hormone deficiency*

Introduction

Children with growth hormone deficiency usually followed up endocrine clinics for treated their short stature and improve their quality of life (1, 2, 3), they aware about their physical growth and feel with embarrassment about body differences from peers 4. As reported in the pediatric studies that children with health disorders may suffer from different negative psychosocial consequences 5. Those children and their parents influence by children's short stature, in addition, to the burden of treatment regime (1, 6). In Baghdad a study conducted in 2009 showed the prevalence of short stature only was 18.7%, while with underweight and short stature was 13.5% 7. The prevalence of short stature related to growth hormone deficiency was 1\4000 child and 60-80% of them with idiopathic causes 17. Many previous literature documented adverse effects

of the short stature on children's psychologically and socially 8.

Methods and Materials:

Research design: cross sectional study used, data was collected from first of May tile the first of November 2018.

Setting: study was carried out in outpatient endocrine clinics at two pediatric teaching hospitals and two specialized centres for endocrine disorders in Baghdad city.

Instrument of the study and procedure: Pediatric Psychosocial Symptoms Inventory PEPSI, parents reported scale, Arabic and English version. It has good reliability, developed from two famous pediatric psychosocial scales: strength and difficulties scale and child behavioral checklist scale by Dr. Al -Ayed and Al-Haider in their study about screening of children's psychosocial problems in Riyadh 9. The approval obtained from the ownerships by electronic mail. PEPSI consists of (38) items, with five subscales: Mood symptoms subscale scored from (0-24), Behavioral

Corresponding Author:

Adraa H. Shawq,

College of Nursing, University of Baghdad,

Email: adraa.hussein@gmail.com

symptoms subscale scored from (0-27), Learning problems subscale scored from (0-18), Somatic symptoms subscale scored from (0-24), and Personality characteristics subscale scored from (0-21). Rating from 0-3: no symptoms= 0, rarely= 1, occasionally= 2, and frequently= 3. The total score ranged from 0-114 (0-38

= mild, 39-76 = moderate, 77-114 = severe).

Statistical Analysis

SPSS programme version 23 was used, frequency, percentage, and mean used in data analysis.

Results



Figure (1) Child's Age and Gender Distribution

In these figures, the mean age of children with growth hormone deficiency is (8.41±2.7), most of them (75%) at school age (6-12) years, and mostly are boys (65%).

Table (1) Pediatric Psychosocial Symptoms Inventory for children with growth hormone deficiency

Levels of scales Frequency	Study Group	
	Percent	
Mild	23	28.75
Moderate	39	48.75
Severe	18	22.5
Total	80	100.0

This table reflects that (84.75%) of children with growth hormone deficiency experience moderate psychosocial problems when assessed by PEPSI.




Figure (2) Subscales Levels of Pediatric Psychosocial Symptoms Inventory for children with growth hormone deficiency

Figure (2) shows the details aspects of psychosocial problems (mood, behaviour, learning, somatic, and personality problems) according to Pediatric Psychosocial Symptoms Inventory PEPSI. Generally, children with growth hormone deficiency experience moderate mood problems (46.6%); moderate behavioural problems (60.4%); moderate learning problems (43.3%) moderate somatic problems (47.6%); moderate personality problems (65%).

Discussion

The results of the present study showed that (48.75% and 22.5%) of children with short stature related to growth hormone deficiency experience moderate and severe psychosocial problems respectively. Their physical growth (height) interferes with social competence and body differences from normal peers which caused psychological disturbances. A study about psychosocial problems of children with growth hormone deficiency by using the Visual Analogue Scale for Children, support the present findings that those children showed negative self perception concerning physical appearance⁶. Another longitudinal study about terminated treatment of adolescents with growth hormone deficiency showed that, their shortness affects negatively on their psychosocial status and their quality of life⁶. Children with growth hormone deficiency suffer from psychosocial disturbances and their psychosocial functioning depending on various factors especially family support and school adaptation¹⁰. Short stature associated with psychosocial distress and emotional deprivation⁵. While another study about reviewing the psychosocial consequences of children with short statured that, those children showed negative social experiences but they have not psychosocial problems⁵. However some studies mentioned the benefit of growth hormone treatment on their quality of life and self esteem after one year of successful treatment⁵.

The present findings showed the aspects of children psychosocial problems according to pediatric psychosocial symptoms inventory PEPSI (learning, somatic, emotional, behavioural, and personality problems), in all aspects of children psychosocial problems children with growth hormone deficiency scored such problems. Mostly they scored behavioural and personality problems (60.4%, 65%). As parent reported their children have mood changes, fight with others, fidgety, tell lies, do not listen to rules, feel down, no hobbies, no friends, spend time alone, depend on

others, and seem shy.

Most of children experienced teasing or juvenalization, these negative psychosocial consequences related to their shortness. Those children experienced behavioral and emotional problems, low self esteem. They exposure to bullying and could not adopt with their condition, the psychosocial burdens may also play role in developing short stature among some children¹¹. Short stature in children can lead to adverse psychosocial impact on those children and affects on their school performance and development of their personality¹. Children with short stature scored behavioural problems as a result of negative social relationships especially with their peers⁶. Another study to measured behavioral and social problems of (195) children with short stature compared with normal children found that children with short stature experience higher score of school difficulties and behavioral problems⁶. Children with short stature experienced different barriers in their personality development which affects negatively on their school and sport competence¹⁷. Children's psychosocial problems and school performance are the most concern of their parent¹¹.

As parent reported children with short stature have low social competence when compared with normal children, because of their unsatisfied appearance which affects on the social relationships and development their personality¹⁰. Those children may face bullying in school or playing with younger children¹⁰. A study in Japan about psychosocial problems of children with growth hormone deficiency by using Child Behaviour Checklist (CBCL) scale, the result of the study showed those children have higher scored than normal children, they also showed high scored on subscale of Child Behaviour Checklist scale such social problems, anxiety and attention⁸. Previous studies showed school and psychosocial problems due to their appearance, they experience internalize problems and behavioral problems, and showed low health related quality of life⁵. By using child behavior checklist scale for children with short stature, the findings showed those children scored high score of behavioral problems than normal developing children in addition to the somatic, attention, and social problems¹⁴. In a study about reviewing children with short stature reported that those children scored lower in academic functioning, while another American study reported no differences form normal children⁵. More than half of children with short stature experienced bullying and showed low quality of life¹⁵.

Children with growth hormone deficiency scored low neuropsychological test and impaired social relation¹³. Those children recorded poor academic level due to effects of social relationships in spite of IQ score comparing with normal children⁸

Kranzler and his colleague in their study about psychosocial function of children with short stature that, those children scored externalize problems and inadequate social relationships¹². Children with growth hormone deficiency showed aggressive behavioral problems, attention, social, emotional problems and maladaptive¹⁵. Prepubertal short children developed more social, internal, and external problems than children in the same age group¹⁰. Children with short stature experience behavioral problems¹⁶. Children with short stature scored low self esteem, bullying, juvenalization, and stigmatization because of their shortness⁵. Previous studies reported those children have academic, internal, external, and behavioural problems due to their shortness⁵. Those children have low quality of life. Psychosocial assessment of children with short stature is important in diagnosis and treatment of those children and well being⁵. Some studies reported those children experienced low quality of life, and internal problems¹⁷. Child health problems affects negatively on their parents.

Early children diagnosis and treatment reduce the risk of psychosocial problems and financial burden of treatment comparing with untreated children, in addition to the psychosocial advantages¹³. Before treatment regimen the psychological profile should be assessed¹⁰. The psychosocial services should also provide during treatment of children with growth hormone deficiency, those children experience sad feelings because of their shortness and puberty development delayed⁴ the psychological assessment of short adult showed some psychological disturbances. The setting and cultural attribute play a role in developing the psychosocial problems associated with certain health problems⁸. The psychological health services will decrease and children well-being improved⁶. Psychosocial improvement is one of the important goals for treated children with short stature¹⁵

Conclusion

Children with growth hormone deficiency were experience psychosocial problems, especially behavioral and personality problems.

Financial Disclosure: There is no financial

disclosure.

Conflict of Interest: None to declare.

Ethical Clearance: All experimental protocols were approved under the College of nursing, University of Baghdad, Iraq and all experiments were carried out in accordance with approved guidelines.

References

1. Jayasena A , Dharshini D , Silva K , Lanka S. Evaluation and management of short stature in children. *Journal of Diabetes Endocrinology and Metabolism* 2011;1
2. Said N, Naji G. Response of Height Measurements to Growth hormone therapy in Short Children with Growth hormone Deficiency. *Iraqi J. Comm. Med.* 2006
3. Al Zubaidi M , Saleh M , Jawad Z. Causes of short stature in Iraqi hospital based study patients, *JFac Med Baghdad* 2017;59: 3.
4. Allen D, MD , Cuttler L. Short Stature in Childhood: Challenges and Choices. *engl j med* 2013;28: 368;13
5. Quitmann JH , Bullinger M , Sommer R , Rohenkohl AC , Silva NM. Associations between Psychological Problems and Quality of Life in Pediatric Short Stature from Patients' and Parents' Perspectives. *PLOS ONE*. 2016.
6. Zlotkin D, Varma SK. Psychosocial Effects of Short Stature. 2006.
7. Al-Saffar AJ. Stunting among primary-school children: a sample from Baghdad, Iraq, *Eastern Mediterranean Health Journal*, 2009;15: 2.
8. Rogol AD. Clinical and Humanistic Aspects of Growth Hormone Deficiency and Growth-Related Disorders. 2012.
9. Al-Ayed IH, Al-Haider FA. Screening For Psychosocial Problems in Children Attending the Pediatric Clinic at King Khalid University Hospital (KKUH) In Riyadh (KSA). *Journal of Family and Community Medicine*. 2008;15(1).
10. Balen HV , Sinnema G , Geenen R. Growing up with idiopathic short stature: psychosocial development and hormone treatment; a critical review. *Arch Dis Child* 2006;91(5): 433–439.
11. Sandberg D E , Colman M , Res H. Growth Hormone Treatment of Short Stature: Status of the

- Quality of Life Rationale. Hormone research in paediatrics. 2005.
12. Kranzler J H , Rosenbloom AL , Proctor B , Diamond FB , Watson M. Is short stature a handicap? A comparison of the psychosocial functioning of referred and non referred children with normal short stature and children with normal stature. *J Pediatr* 200;136(1):96-102.
 13. Stagi S , Scalini P , Farello G , Verrotti A. Possible effects of an early diagnosis and treatment in patients with growth hormone deficiency: the state of art. *Ital J Pediatr* 2017; 43: 81. Published online 2017 Sep 16.
 14. Stabler B , Siegel PT , Clopper RR , Stoppani C E , Compton PG , Underwood L E. Behaviour change after growth hormone treatment of children with short stature. *J Pediatr*. 1998;133(3).
 15. Tanaka T , Matsuo N , Seino Y , Fujieda K , Yokoya S , Tachibana K.,the Study Group of GH Treatment in IUGR. Growth-Promoting and Psychological Effects of High-Dose Growth Hormone Treatment in Children with Intrauterine Growth Retardation. *Clin Pediatr Endocrino* 2000;9(1): 7-17.
 16. Kremidas D, Wisniewski T, Divino V M, Bala K, Olsen M, Germak J, Aagren M, Holot N, Lee W. Administration Burden Associated With Recombinant Human Growth Hormone Treatment: Perspectives Of Patients And Caregivers. *Journal of pediatric nursing* 2013;28: 55-63
 17. Silva N, Bullinger M, Sommer R, Rohenkoh A, Witt S, Quitmann J. Children`S Psychosocial Functioning and Parent`S Quality Of Life in Paediatric Short Stature: The Mediating Role of Caregiving Stress. *Clin psychol psychother.* 2017;21-12.