

The Comparison of Social Support Against the Life Quality of The Spinal Cord Injury Under Stress

Soudabeh Aloustani¹, Karvan Bekmaz², Adeleh Sadeghloo³, Hamid Hojjati¹

¹Lecturer, Department of Nursing, Aliabad Katoul Branch, Islamic Azad University, Aliabad Katoul, Iran,

²Lecturer, Department of Nursing, Khoy Branch, Islamic Azad University, Khoy, Iran, ³ lecturer, Young Researchers and Elite Club, Aliabad Katoul branch, Islamic Azad University, Aliabad Katoul, Iran

Abstract

Introduction: The stress of war has caused severe problems for the individual and family. Spouses of veterans are gradually involved in family erosion. Lack of social protection and social dysfunction decreased life satisfaction and reduce intimacy between partners.

Purpose: this study has been done in the purpose of comparing social support against the veteran's life who are under stress.

Method: This cross-sectional study was done on 79 posttraumatic stress disorder veteran wives and it was also done on 79 ordinary wives through convenience sampling. The instrument to collect the required information was the quality of life questionnaire which contained 36 SF questions and Medical Outcomes Study Social Support Survey (MOS-SS).and the needed information was collected by self-reported method.

Results: Veteran wives social support with the mean and the standard deviation was 19.3 ± 5.1 while it was 31.1 ± 3.7 versus the ordinary wives and this difference was significant (Pless than 0.01).It showed the veteran wives quality of life with the mean and standard deviation of 46.3 ± 0.088 while it showed 66.9 ± 8.5 the mean and standard deviation for the ordinary wives and it showed the significant difference (Pless than 0.01).

Conclusions: the obtained result showed that the quality of life related to the health and social support in veteran wives is lower in compare with the ordinary wives with the same situation, therefore educational, treatment and supporting and promoting plans for this group is necessary.

Key word: social support; life quality; veterans; PTSD

Introduction

It is important to pay attention to mental health^(1, 2). These effects include spinal cord injury⁽³⁾. Chronic diseases threaten the physical and mental health of the patient⁽⁴⁾. Posttraumatic stress disorder is a serious reaction and perpetual response which appears because of being exposed to serious stresses⁽⁵⁾. Posttraumatic stress disorder was identified and recognized during the civil war in the United States⁽⁶⁾. The prevalence of this

disorder in the public population is 4-8 percent⁽⁷⁾. But this percentage is between 9 to 25 percent among the war injured ones during the first two years⁽⁸⁾ The prevalence of that among the Vietnamese war soldiers has been reported as much as 30.1% to 14.7%⁽⁹⁾. It has also been reported 17.1% to 15.6% for the solders coming back from Iraq war^(10, 11) According to the present documents more of the Iranian veterans have posttraumatic stress disorder and this disorder which is always with the veterans has a very bad effect on they themselves, their lives and their family lives^(12, 13) Living with a person who suffers PTSD can transfer some of the symptoms to their wives⁽¹⁴⁾.so one of the war victims is the veteran wives who are exposed to the PTSD so the veteran wives encounter lots of problems in their lives and theses

Corresponding author:

Hamid Hojjati

lecturer, Department of Nursing, Aliabad Katoul Branch, Islamic Azad University, Aliabad Katoul, Iran
E-mail: hojjati_hamid2019@yahoo.com

problems become more and more by passing the time and cause them to have mental^(15, 16) and this continual disorder can bring trauma to the wives and can cause anxiety, depression, isolation in the veteran wives^(17, 18). The veteran wives will have family erosion little by little and because of the lack of social support they lose their satisfaction with the family members especially with their husbands⁽¹⁹⁾. One of the anxiety of the family suffering from PTSD is mental relationship, matrimony compatibility and sexuality satisfaction which can affect their life quality⁽¹²⁾. Most PTSD veteran wives sometimes face challenges that force them to be compatible to life because of their children and loving their husbands and some wives who cannot tolerate this hard situation decide to divorce⁽²⁰⁾. Studies have shown that most veteran wives have lack of social communications and they cannot express themselves well and they hesitate whether to continue that life or separate their husbands and in most cases the presence of their children cause them not to do so and this causes them to accept the problems and become compatible⁽²¹⁾. An eight-year war with Iraq and its follow-ups have made many Iranian warfare Died and veterans suffer from war-related Mental disorders⁽²²⁾. In addition to medical and Mental disorders, this disorder decreases the sufferers' quality of life⁽¹⁶⁾. So, this study that has been based on the purpose of comparing the social support and life quality of PTSD veteran wives has been done.

Material and Method

This cross-sectional study was done on two wife groups; the first group were the veteran wives and this group contained 158 PTSD veteran wives and the second group consists of 79 ordinary wives. This study was done in the year 2016. The study setting was Aliabad city, the population of the city is 150,000 people and according to the announcement of The Foundation of Martyrs and Veteran Affairs, there are 96 PTSD veterans there. The method of sampling was done by census method among the ones who had inclusion criteria which was certified by a psychologist, and those samples were the veterans with documents that proved they are veterans, and not addicted to any drug and abuse substances were chosen. All participants provided informed consent after the study purpose was explained, and assurances were made of anonymity and privacy. At most, 79 people were qualified for the study. The control group was elected among the veteran wife relatives and their close friends and they had no PTSD disorder and no chronic mental disorder and they were not addicted to any drugs.

This study information was collected by Demographic questionnaire, Medical Outcome Study Social and supportive Survey (MOS-SS).

Demographic questionnaire: This questionnaire contains information such as age, education, number of children, the length of time living with his wife, drug using, and the chronic mental illnesses.

Short form of quality of life (QOL-SF36): This questionnaire which contains 36 questions measures and evaluates the physical, mental and eight domains of life. The physical life quality consists of physical function, limitation in having a role because of the physical lack in efficiency, physical pain and general health.; and domain of psychological, life quality, contains limitation in role because of having emotional problems, social function, happiness and vividness and healthy mental. The method of giving score to them was between zero for the worst to 100 according to the questionnaire instruction SF36, and the score obtained by the samples is between 0 and 100 and the reliability of this study has strongly been confirmed and the reliability of the native form of this questionnaire by Shahgholian(2014) in all aspects has firmly been confirmed and is 0.95 – 0.91⁽²³⁾.

Medical Outcomes Study Social Supports Survey (MOS-SS): This questionnaire was made by Sherbourne and Stewart. These supportive social tools have 19 questions and 5 sub scales. These subscales are tangible support (4 questions), which measures the financial and behavioral, exciting support (4 questions) which evaluates the positive emotion, sympathy, encouragement to express the feelings. Informing (4 questions) which evaluates guidance, informing, kindness (3 questions) which evaluate the amount of love, and positive social compatibility (4 questions) which evaluates the people who do the amusement and fun activities. This scale is a self-report tool which is used to measure the amount of agreement or disagreement of someone to any of them in five-point Likert scale (never = 1 point, seldom = 2 points, sometimes = 3 points, often = 4 points, always = 5 points). The lowest score in this is 19 and the highest score is 95. The high point in this test shows that the test is supported by the society a lot⁽²⁴⁾. The reliability of sub domain of this test was reported by Cronbach's alpha 0.91⁽²⁵⁾. All statistical procedures were performed with SPSS 16 statistical software. Using descriptive statistics independent t-test.

Findings

The mean age of the PTSD veteran wives were 39 individuals (49%), high school and guidance school education was 24 individuals (31%) primary school education 13 individuals (6%) and three of them had university degree, among the ordinary people wife, 48 (61%) had high school and guidance school education, (31%), 5 individuals were uneducated, 5 individuals (6%) had diploma and two of them (3%) had university degree.

The amount of life quality of the veteran wives with means and standard deviation was 46.3 ± 0.88 ($P < 0.01$). There also was seen a significant difference between the veteran wives versus the ordinary people (Table 1).

The amount of veteran wife’s social support by mean and standard deviation was 19.3 ± 5.1 and in the ordinary people it was 31.1 ± 3.7 ($P < 0.1$).

Table (1): comparing the PTSD veteran wife’s life quality

Dimension of the group	Life quality		P_value
	Veteran wives	Ordinary people wives	
Physical dimension	24.1 ± 4.9	31.4 ± 4	$p < 0.01$
Mental dimension	22.2 ± 5.5	35.5 ± 4	$p < 0.01$
Life quality general score	46.3 ± 8.8	66.9 ± 8.5	$p < 0.01$

Discussion

Spinal cord injury affects the mental health of patients, with a prevalence of mild depression of 22.6% in the Hatefi et al⁽²⁶⁾. The result of the study shows that the PTSD veteran wives have a lower quality of life. In both the physical and mental dimensions, the veteran wives had lower quality of life in compare with the ordinary people wives. Studies have shown that the veteran wives have more excitement and it has caused them to have a lower quality of life. The results of the present study were same as Bares (2011), Brow (2010) and in all these studies the PTSD veteran wife’s life quality was low^(9, 27).

Clinical experiences have shown that the veteran family, particularly their wives, suffer mental problems such as seclusion feeling, loneliness, inability, sin feeling, and depression⁽²⁸⁾. because the PTSD veteran wives are exposed to stresses and have lots of turmoil in their marriage lives⁽¹⁵⁾. Lots of studies show that the PTSD veteran wives suffer a lot from mental disorder in compare with the ordinary people or the wives of the veteran who have no stress^(16, 29, 30). shows the surveys of this toll in the form of depression, anxiety and mental

disorder and also as the physical problems and it is known as the prevalent headache and physical complaint among the veteran wives.²⁷ because living for a long time with someone who suffers post-traumatic stress disorder has a deep and serious influence on their family and can be a chronic stress factor for their wives and can be the source of big and serious problems their families⁽³¹⁾.

This study showed that the social support for the veteran wives is lower than ordinary people wives. Communication disorder, physical problems and its consequence, lack of understanding the veterans by some people in the society is one of the most important things that the veterans suffer from⁽¹⁵⁾. Social supports may have a very important effect on developing care health and also on mental pathology in reaction to the stressful life events⁽³²⁾. Social supports from the authorities and the social organizations can increase respect, can reduce blame and label and it can also form a better self-compatibility in veteran wives and it also can reduce stress in their lives⁽¹⁶⁾.

Conclusion

According to the findings of this study, it can

be concluded that the martyr and veteran families are special families with special stresses and the amount of their stress increases day by day. Women are the center of these stresses, and they themselves and their children are at the risk of danger therefore the society is at danger. Therefore, in order to reduce the amount of stress and the factors that causes stress and anxiety in this group, presenting more and more suitable services and economical is an evitable necessity. Besides, in order to reduce the damage of stresses, teaching educational services, that is teaching life skills are among the communicative skills and stress management and doing actions on beliefs and their attitude towards the incidents and life meaning will be useful. It is also suggested that the veteran wife's life compatibility is to be investigated through a quantitative research.

Conflict of Interest: There is no conflict of interest between authors.

Acknowledgment: This article is the result of Research Project No. 94413 Approved by the Young Researchers Club, Islamic Azad University of Aliabad Katoul. We appreciate the cooperation of the Foundation of Martyrs and Veterans Affairs and the participants in conducting this research.

Source of Funding: This work was supported by the Young Researchers Club, Islamic Azad University of Aliabad Katoul.

Ethical Clearance: Ethical considerations in this study included obtaining written consent from participants, assuring them of the confidentiality of their data, and allowing them to freely participate in this research.

Reference

1. Karimian M , Borji M, Otaghi M, Mansouri F, Solaimanizadeh L, et al. Special Attention to Psychiatric Centers in COVID-19 Pandemic, Arch Clin Infect Dis. Online ahead of Print ; 15 (2):e103258. doi: 10.5812/archcid.103258.
2. Karimian M, Mansouri F, gheiasi GH, Psychological Abuse in Elderly during Exchanging News of COVID-19 , Iranian Journal of Psychiatry and Behavioral Sciences, 2020, 13 (3). 22-24.
3. Vaccarino V, Goldberg J, Rooks C, Shah AJ, Veledar E, Faber TL, et al. Post-traumatic stress disorder and incidence of coronary heart disease: a twin study. 2013;62(11):970-8.
4. Otaghi M, Shohani M, Sayehmiri K, Razaghi MJJoFM, Toxicology. The effect of orem's self-care program on adherence to treatment among the elderly patients with type II diabetes in ilam in 2018. 2018;12(4):189-94.
5. Vaccarino V, Goldberg J, Rooks C, Shah AJ, Veledar E, Faber TL, et al. Post-traumatic stress disorder and incidence of coronary heart disease: a twin study. 2013;62(11):970-8.
6. Crocq M-A, Crocq LJDien. From shell shock and war neurosis to posttraumatic stress disorder: a history of psychotraumatology. 2000;2(1):47.
7. Widome R, Laska MN, Gulden A, Fu SS, Lust KJAJHP. Health risk behaviors of Afghanistan and Iraq war veterans attending college. 2011;26(2):101-8.
8. Dohrenwend BP, Turner JB, Turse NA, Adams BG, Koenen KC, Marshall RJS. The psychological risks of Vietnam for US veterans: a revisit with new data and methods. 2006;313(5789):979-82.
9. Braš M, Milunović V, Boban M, Brajković L, Benković V, Đorđević V, et al. Quality of life in Croatian Homeland war (1991-1995) veterans who suffer from post-traumatic stress disorder and chronic pain. 2011;9(1):56.
10. Connell M, Omole O, Subramaney U, Olorunju SJAjop. Post traumatic stress disorder and resilience in veterans who served in the South African border war. 2013;16(6).
11. Friedman MJ. Acknowledging the psychiatric cost of war. Mass Medical Soc; 2004.
12. Klarić M, Frančišković T, Pernar M, Nemčić Moro I, Miličević R, Černi Obrdalj E, et al. Caregiver burden and burnout in partners of war veterans with post-traumatic stress disorder. 2010;34(1):15-21.
13. Evans L, McHugh T, Hopwood M, Watt CJA, Psychiatry NZJo. Chronic posttraumatic stress disorder and family functioning of Vietnam veterans and their partners. 2003;37(6):765-72.
14. Kirby AC, Beckham JC, Calhoun PS, Roberts ST, Taft CT, Elbogen EB, et al. An examination of general aggression and intimate partner violence in women with posttraumatic stress disorder. 2012;27(5):777-92.
15. Klarić M, Frančišković T, Černi Obrdalj E, Petrić D, Britvić D, Zovko NJPD. Psychiatric and health impact of primary and secondary traumatization in wives of veterans with posttraumatic stress

- disorder. 2012;24(3.):280-6.
16. Peraica T, Vidović A, Petrović ZK, Kozarić-Kovačić DJH, outcomes qol. Quality of life of Croatian veterans' wives and veterans with posttraumatic stress disorder. 2014;12(1):136.
 17. Henry SB, Smith DB, Archuleta KL, Sanders-Hahs E, Goff BSN, Reisbig AM, et al. Trauma and couples: Mechanisms in dyadic functioning. 2011;37(3):319-32.
 18. Tsai J, Harpaz-Rotem I, Pietrzak RH, Southwick SMJPI, Processes B. The role of coping, resilience, and social support in mediating the relation between PTSD and social functioning in veterans returning from Iraq and Afghanistan. 2012;75(2):135-49.
 19. Frančišković T, Stevanović A, Jelušić I, Roganović B, Klarić M, Grković JJCmj. Secondary traumatization of wives of war veterans with posttraumatic stress disorder. 2007;48(2.):0-184.
 20. Mansfield AJ, Williams J, Hourani LL, Babeu LAJJoTSOPoTISfTSS. Measurement invariance of posttraumatic stress disorder symptoms among US military personnel. 2010;23(1):91-9.
 21. Lyons MAJJoAN. Living with post-traumatic stress disorder: the wives'/female partners' perspective. 2001;34(1):69-77.
 22. Azad EM, Hashemi SZJjop. The Effectiveness of Mindfulness Training in Improving the Quality of Life of the War Victims with Post Traumatic stress disorder (PTSD). 2014;9(4):228-36.
 23. Shahgholian N, Eshghinezhad A, Mortazavi MJJjon, research m. The effect of tai chi exercise on quality of life in hemodialysis patients. 2014;19(2):152.
 24. Moser A, Stuck AE, Silliman RA, Ganz PA, Clough-Gorr KM. The eight-item modified Medical Outcomes Study Social Support Survey: psychometric evaluation showed excellent performance. *Journal of clinical epidemiology.* 2012;65(10):1107-16.
 25. Shokrvash B, Salehi L, Akbari MH, Mamagani ME, Nedjat S, Asghari M, et al. Social support and dairy products intake among adolescents: a study from Iran. 2015;15(1):1078.
 26. Hatefi M, Abdi A, Tarjoman A, Borji MJTM. Prevalence of depression and pain among patients with spinal cord injury in Iran: A systematic review and meta-analysis. 2019;24(4):e87503-e.
 27. Bravo-Mehmedbašić A, Kučukalić A, Džubur Kulenović A, Suljić EJPd. Impact of chronic posttraumatic stress disorder on the quality of life of war survivors. 2010;22(3):430-5.
 28. Vagharseyyedin SAJPHN. Experiences of wives of Iranian war veterans with posttraumatic stress disorder regarding social relationships. 2015;32(2):122-31.
 29. Caska CM, Renshaw KDJJoAD. Perceived burden in spouses of National Guard/Reserve service members deployed during Operations Enduring and Iraqi Freedom. 2011;25(3):346-51.
 30. Renshaw KD, Caska CMJBT. Relationship distress in partners of combat veterans: The role of partners' perceptions of posttraumatic stress symptoms. 2012;43(2):416-26.
 31. Galovski T, Lyons JAJA, behavior v. Psychological sequelae of combat violence: A review of the impact of PTSD on the veteran's family and possible interventions. 2004;9(5):477-501.
 32. Shyu Y-IL, Tang W-R, Liang J, Weng L-JJNr. Psychometric testing of the social support survey on a Taiwanese sample. 2006;55(6):411-7.