

Optimization of Streptokinase Mutant Protein Purification Method Using Affinity Chromatography Technique

Nastaran Rahimi¹, Mohammad AlinezhadChamazketi², Amir Yaghoubi Nezhad³, Forough Talaeizadeh⁴

¹Research assistant in the Islamic Azad University Pharmaceutical Sciences Branch, Tehran, Iran, ²Lecturer in Department of Biotechnology, Malek Ashtar University of Technology (MUT), Tehran, Iran, ³Department of Cellular and Molecular Biology, Faculty of Sciences, Higher Education Institute of Rab-Rashid, Tabriz, Iran, ⁴Department of Educational Science and Psychology, Islamic Azad University Marvdasht Branch, Marvdasht, Iran

Abstract

Protein purification has always been one of the most critical and challenging stages of drug-protein production. Streptokinase as the most common, and currently, the most cost-effective fibrinolytic drug is no exception. In this study, the mutated streptokinase producing clone (SK263cyc) to which the histidine tag was added was grown in TY2x medium, and SDS-PAGE assessed protein expression after induction of protein expression. Three different methods did protein purification; in the first one, metal, ion affinity chromatography (IMAC) technique was used. In the second solution, first, by filtration with ammonium sulfate, the purification was carried out, and then by affinity purification, chromatography continued. In the third solution, hydrophobic chromatography was used to purify the streptokinase protein. The purity of ophthalmic purity was 93.2%, and the purity of hydrophobic purity was about 90.4%, whereas the combination of pre-treatment with ammonium sulfate and the purity of the ophthalmic method did not achieve more than 88%. In general, the results of this study show that the IMAC method is more suitable as a final method in the process of streptokinase purification than the other two approaches.

Keywords: Streptokinase, Recombinant protein, Affinity chromatography, Hydrophobic gel purification, Ammonium sulphate

Introduction

When a blood clot forms in the circulatory system, it can cause blood clots, which can cause dangerous complications and, ultimately, death. A healthy balance system prevents clots from forming in a natural circulation system⁸. The mechanism of blood clot formation is one of the most important and complex physiological systems in the body, and streptokinase is the most common and currently the most cost-effective fibrinolytic drug for this purpose¹⁻⁸. The most important part of designing a suitable “downstream process” for purification of a protein for analysis and treatment purposes is the separation of the desired protein from other available impurities and contaminants using various purification methods⁹⁻¹¹. The highest increase in concentration is due to the separation stage of the product, but the quality of the products dramatically increases during the initial purification stage^{3, 12-16}. SK263cys is a mutant protein that contains cysteine amino acid at position 263 instead of glutamic acid and is designed to

perform specific pegylation on cysteine amino acid. High purity protein is required for this process. Generally, in industrial processes and most research, the HIC method is used to purify streptokinase. Because this process is costly and low yield¹⁷⁻¹⁸. In this study, purification of streptokinase protein was performed by applying and optimizing the IMAC method and combining it with one step pre-purification by ammonium sulfate deposition method. We also tried to purify this protein with the highest purity and higher efficiency^{8, 11}.

Materials and Method

Protein Expression

The most commonly used host for the expression of Escherichia coli, the strain of Escherichia coli BL21 Rosetta (DE3), was used to express the mutant streptokinase protein. First, in a 5cc TY2x medium containing kanamycin antibiotic 50 ml / µg, a unique culture was prepared overnight. After 24 hours, after transferring the vertex tube, 100 µl of this medium was

transferred to 50 ml of fresh and warm TY2x medium containing kanamycin antibiotic and stirred at 150 rpm at 37 ° C until ambient light absorption at 600 nm (600 OD) (0.4 to 0.6).

To produce protein and induce protein expression in large quantities, 50 µl of stock (M1 IPTG) was added to the culture medium to increase its concentration in 1 mM medium. The medium environment-friendly was returned to the incubator again and incubated for 3–4 h at 150 rpm at 37 ° C. After incubation, the OD600 medium was re-recorded.

Protein Purification

Cell failure was performed by cell lysis (8 mM urea + 10 mM tris + 100 mM sodium dihydrogen phosphate). To prepare cell extracts from 50 ml TY2x culture medium, the culture medium was pre-divided into two falcons and centrifuged at 5000 rpm for 10 min.

The supernatant was discarded, and the precipitate was kept. To the precipitate was added 5 ml of lubricating buffer. After one to two hours, the crystalline contents of the Falcon were centrifuged at 5000 rpm for 10 minutes, and the supernatant was kept in the freezer after separation. The cell extract obtained was used in three strategies.

- 1: Hydrophobic Interaction Chromatography
- 2: Immobilized Metal Affinity Chromatography

The first strategy is the use of affinity chromatography without pre-purification, and the second strategy is the use of ammonium sulfate precipitation as the pre-purification and the affinity chromatography as the final purification. Both of these solutions were compared with the primary approach used in the industry.

Purification by Hydrophobic Interaction Method (HIC)

Gel filtration chromatography was performed using Sephadex-25G gel. By passing column equilibration buffer (20 mM tris, 0.2 mM sodium chloride and 8.5 = pH), the sample was prepared for injection, and then 5 mL of sample was slowly loaded into the column by the sampler.

Purification by Affinity Chromatography

One of the important features of this Purification

system is its high capacity for binding to His-tag proteins (5-10 ml / mg) and insensitivity of the system to the precise 3D structure of the protein. For small-scale denaturing purification, after breaking the cell wall and extracting the cell contents, we centrifuge the liquid and load the supernatant after centrifugation on the chromatography column. After complete removal of buffer (pH 8), urea 8 mM + 10 mM tris + 100 mM sodium dihydrogen phosphate (5 ml sample was injected into the column by the sampler). When the sample was completely removed from the column, the washing buffer passed through the column. In the process of this study, pH of the buffer of the washing step was optimized in the range of 6.3 to 5.3 and a suitable pH of 5.7 was obtained. Figure 2b shows the effect of pH regulation on the efficiency of the IMAC method. At this point, proteins that are weakly bonded with nickel or have no bond at all with histidine are removed from the column. If the pH of the buffer used in the washout phase is not adequate, the target protein may also be removed from the column, resulting in a negative effect on the final purification. In the wash step, the buffer is introduced into the column at pH 4.5 to remove the target protein. The column outlet was collected in sterilized microtubes in a volume of 1 ml [4, 10].

2.2.3 Pre-Purification by Ammonium Sulfate Deposition and Purification by Affinity Chromatography:

After preparation of the cell extract, the product was prepared with a pre-purification step by ammonium sulfate precipitation to enter the IMAC column. The amount of protein in the samples from each step of the process was determined by electrophoresis. Figure 5 illustrates these results. Table 3 also relates to the results of the gel analysis by software.

Protein measurement

Method: SDS-PAGE

To confirm the presence of the protein and to evaluate its value, the samples obtained from each step of the studied strategies were used by PAGE-SDS electrophoresis [7].

Optical Method for Calculating Total Protein:

OD of each sample was measured using a spectrophotometer at 280 nm. The mean ODs were obtained and multiplied by the dilution and extinction coefficient of 0.925 for streptokinase, and the total protein content of the sample was obtained [9].

Calculate the process efficiency for the target protein and its purity

The gel electrophoresis image of each process was analyzed by Lab Image software, and the percentage of protein purity was determined at each step; the purification efficiency is then calculated using the degree of purity. degree of purity = Total Streptokinase / Total Protein Concentration * 100

Yield = Purified Protein Concentration * Degree of purity / total protein concentration

Results

The results of electrophoresis showed that the protein was produced properly (Fig. 2). In the HIC purification strategy, the urea in the protein solution was first removed (Fig. 1A), and then the isolated proteins were isolated from streptokinase by HIC using S-650 butyl gel (Fig. 1B).




Figure 1A. Chromatogram obtained by gel filtration chromatography; B. Chromatogram obtained by HIC.

In vitro analysis of purified samples showed that all three purification strategies performed well. After optimizing the pH of the wash buffer, the amount of protein produced by IMAC at a concentration of 1.71 mg / ml was higher than the other two methods (Fig. 2B). After careful examination using Lab Image software and determination of purity, it was found that the highest purity was related to IMAC93% method. The results of electrophoresis showed that the protein was produced at the appropriate level at the beginning of the work.




Figure 2. Electrophoresis of the products of the investigated methods. A shows the electrophoresis of purified products by HIC method. L3: protein before purification, L4: sample after gel filtration step, L5: Sample after ammonium sulfate deposition, L6: sample after the HIC step b. The results of electrophoresis of the products show the method (IMAC method 1). L2: sample before purification, L4: (Sample 1) Wash buffer purification (pH = 6.3), L5: sample (2) Purification by washing buffer pH = 6, L6: Sample 3 (Purification by washing buffer, L7: (pH = 5/7) Sample (4) purification by washing buffer (pH = 5/3 C. Results of electrophoresis of the products of pre-purification by ammonium sulfate precipitation and purification by IMAC method) shows the two columns in the image, L3: protein before purification, L4: Sedimentation step protein, L5: First output, L6: IMAC Second output IMAC.

Measurement of protein concentration by the HIC method showed that the highest amount of protein and the highest yield was obtained by the IMAC method, while the lowest yield was obtained by combined IMAC method and deposition with ammonium sulfate (Fig. 3A and B).




Figure 3. Comparison of Purification Results with Different Methods; A - Comparison of Protein Levels with Three Purification Methods; B-Comparison Chart and Purity Efficiency with Three Purification Methods.

Conclusion

The results of this study showed that the IMAC method had better efficiency than HIC in purification of streptokinase with histidine tag. Also during the IMAC process based on the results, it was found that the pH of the wash buffer for the purification of streptokinase protein was 5.7.

Ethical Clearance: This project didn't need any ethical permission.

Source of Funding: The project was co-funded through Islamic Azad University Pharmaceutical Sciences Branch, Tehran, Iran and Iranian-Australian Community of Science, Tehran, Iran.

Conflict of Interest: There is no conflict of interests in this project.

Reference:

- [1] Mehrabifar A, Rahmati M. Significant Increasing of Gastrointestinal Dissolution of Aceclofenac via Florite. *Medbiotech Journal*. 2017;01(01):9-14.
- [2] Hochuli, E. Bannwarth, Dobeli, W. H. Gentz and Suber, D., Genetic approach to facilitate purification of recombinant proteins with a novel metal chelate adsorbent. *Biotechnology*. 1998, 6: 1321-1325.
- [3] Zendegany S. Social and civil perceived supports for kidney patients compared to normal people. *Journal of Humanities Insights*. 2019;3(01):17-9.

- [4] H. Block, B. Maertens, et al., 2009. Immobilized-Metal Affinity Chromatography (IMAC); Methods in Enzymology, Volume 463; Elsevier Inc.
- [5] Farzadnia E, Hosseini Z, Riahi A. Study of Hospital Quality Management and Improvement Rates in the Hospitals. *Journal of Humanities Insights*. 2017;01(01):7-11.
- [6] Jafari R, Mirshahi M, Production NC, van Es T and Davis FF, 2007, Effect of covalent attachment of polyethylene and purification of recombinant streptokinase using pMAL expression vector, *Tehran University Medical Journal*; Vol. 65, No. 1
- [7] Wat P. Pb and Cd Detection in some Pharmaceutical Products through Electrochemical Method. *Medbiotech Journal*. 2017;01(01):22-8.
- [8] Monzavi et al., Design, cloning, Expression and evaluation of cysteine-substitutes of intact and truncated molecules of streptokinase, *Physiology and Pharmacology*, Spring 2010, 14 (1), 56 – 65.
- [9] Barbosa F. Emergency Health Care System and its Role in National Disasters. *Journal of Humanities Insights*. 2018;02(01):14-20.
- [10] Farzadnia E, Hosseini Z, Riahi A. Study of Hospital Quality Management and Improvement Rates in the Hospitals. *Journal of Humanities Insights*. 2017;01(01):7-11.
- [11] Amanlou M, Mostafavi SM. In silico screening to aim computational efficient inhibitors of caspase-9 by ligand-based pharmacophore modeling. *Medbiotech Journal*. 2017;01(01):34-41.
- [12] Bouaffad H, Hayyani N. Determination of Elemental Impurities Ciprofloxacin hydrochloride and Fluconazole via ICP-OES. *Medbiotech Journal*. 2018;02(01):41-6.
- [13] Heidari S, Imani M, Mostafavi SM. A Validated and Rapid HPLC Method for Quantification of Human Serum Albumin in Interferon beta-1a Biopharmaceutical Formulation. *MedBioTech Journal*. 2017;1(01):29.
- [14] Heidary S, Imani M, Mostafavi SM. A Validated and Rapid HPLC Method for Quantification of Human Serum Albumin in Interferon beta-1a Biopharmaceutical Formulation. *Medbiotech Journal*. 2017;01(01):29-33.
- [15] Huang, C. J., Lin, H., and Yang, X., Industrial production of recombinant therapeutics in *Escherichia coli* and its recent advancements. *Journal of industrial microbiology & biotechnology*, 2012, 39(3): p. 383-399
- [16] Mostafavi SM, Bagherzadeh K, Amanlou M. A new attempt to introduce efficient inhibitors for Caspas-9 according to structure-based Pharmacophore Screening strategy and Molecular Dynamics Simulations. *Medbiotech Journal*. 2017;01(01):1-8.
- [17] Obidoa P, Ugueche U. In-vitro Investigation of Ocular E-Coli Infection via Norfloxacin. *Medbiotech Journal*. 2017;01(02):96-103.
- [18] Samira Eissazadeh MP, Mohammad Sadegh Taskhiri, Mostafavi SM. Improvement of Sensitivity of Antigen-Antibody Detection of Semen Through Gold Nanoparticle. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 2019;10(1):144.