

Incidence of BCC in a Tertiary Care Hospital in Malwa region of Punjab

Grace Budhiraja¹, Pulkit Bharti², Harsimrat Singh³

¹Professor, ²Post Graduate, ³Senior Resident, Department of Ent, Adesh Institute Of Medical Sciences And Research, Bathinda, Punjab

Abstract

Objectives. Although the incidence of skin cancers in India (part of South Asia) is low, the absolute number of cases may be significant due to large population. The existing literature on BCC in India is scant. So, this study was done focusing on its epidemiology, risk factors, and clinicopathological aspects.

Methods. A hospital based cross-sectional study was conducted in Punjab, North India, from 2017 to 2019. History, examination and histopathological confirmation were done in all the patients visiting skin department with suspected lesions.

Results. Out of 36 confirmed cases, 63.9% were females with mean \pm SD age being years. Mean duration of disease was 4.7 years. Though there was statistically significant higher sun exposure in males compared to females (value being 0.000), BCC was commoner in females, explainable by intermittent sun exposure (during household work in the open kitchens) in women. Majority of patients (88.9%) had a single lesion. Head and neck region was involved in 97.2% of cases, with nose being the commonest site (50%) with nodular/noduloulcerative morphology in 77.8% of cases. Pigmentation was evident in 22.2% of cases clinically. Nodular variety was the commonest histopathological variant (77.8%).

Conclusions. This study highlights a paradoxically increasing trend of BCC with female preponderance, preferential involvement of nose, and higher percentage of pigmentation in Indians.

Keywords: basal cell carcinoma; pigmentation; histopathological

Introduction

Jacob Arthurin 1827 first coined the term “rodent ulcer” to describe what we now know as a basal cell carcinoma (BCC) [1]. It is the most common cutaneous malignancy worldwide, accounting for 65–75% of all skin cancers. Gross differences are noted in the percentage of skin cancer in the Asians (2–4%) and Blacks (1-2%) as compared to the Caucasians (35–40%) [2]. Although the incidence of skin cancers in India is lower as compared to the Western world, absolute number of cases may

be significant due to large population. The existing literature on BCC in India is scant with lack of clinical studies with statistical analysis [3]. So, this study was undertaken to fill this deficit in literature of BCC with focus on epidemiology, risk factors, and clinical and pathological aspects of the disease.

BCC is a nonmelanocytic skin malignancy arising from basal cells of the epidermis or follicular structures and is seen mostly on sun exposed areas, especially head and neck, occasionally over the trunk and limbs, and rarely on the palms, soles, mucous membranes, and genitals [4, 5].

The anatomic distribution of BCC correlates with embryonic fusion planes. Recently, it has been indicated that BCC occurrence is higher along embryonic fusion planes as compared to other areas of the midface, evidence

Corresponding Author:

DR. Pulkit Bharti

Post graduate, Department of ENT
Adesh institute of medical sciences
Pulkitbharti58@gmail.com, 9876319803

that supports this hypothesis for BCC pathogenesis [6]. (SD), and frequency tables were utilized.

Ninety-five percent of these neoplasms occur in patients aged more than 40 years, although cases in childhood and congenital basal cell epitheliomas have been reported [7-9]. In children, it is usually associated with a genetic defect, such as basal cell nevus syndrome, xeroderma pigmentosum, nevus sebaceous, epidermodysplasia verruciformis, Rombo syndrome, or Bazex syndrome.

Material and Methods

A hospital based study was conducted at a tertiary care hospital situated in Punjab, North India, from 2017 to 2019. Patients of all ages attending skin outpatient department with suspected lesions were screened for BCC after taking an informed written consent. Patients with histopathologically confirmed BCC were enrolled in the study.

Detailed history with recording of various patient variables like age, gender, duration of symptoms, Fitzpatrick skin phototype, skin color, average daily sun exposure (hours/day), occupation, residence place (rural or urban), exposure to chemicals including pesticides, radiation exposure history, treatment with psoralen UVA (PUVA) or narrow band UVB (NBUVB), smoking, alcohol intake, history of personal or family history of skin cancers, personal or family history of other cancers, history of genetic disorder like xeroderma pigmentosum, albinism, and history of previous treatment.

Clinical examination was done with data collection on various tumor variables which included the following: size, location, number, morphological subtype, and pigmentation. For descriptive purposes, the lesions were classified based on size into small (less than 1 cm in diameter), medium (1-2 cm in diameter), and large (>2 cm in diameter).

Investigations included complete blood count with differentials, bleeding time, clotting time, renal function tests, liver function tests, and viral markers. Additional investigations were done depending upon the clinical scenario. Diagnosis was confirmed by histopathological examination of biopsy specimen with documentation of histopathological variant. To analyze the results, descriptive statistics such as mean, standard deviation

Results

A total of 36 histopathologically confirmed cases of BCC were enrolled in the study from 2011 to 2013. An increase was seen in absolute number of cases diagnosed per year with 9, 11, and 16 patients in 2017, 2018, and 2019, respectively .

Out of these patients, males were 36.1% (13/36) and females were 63.9% (23/36) with M:F being equal to 0.57:1 . Age of the affected cases ranged from 29 to 92 years of age. The mean \pm SD age of the patients was years (60.9 yrs years for males and 57.6 years in case of females). Although the difference in mean age between males and females was not statistically significant (data was analyzed using unpaired -test), it carries a clinical relevance as females tend to seek medical care earlier than males for suspicious, asymptomatic, and cosmetically disfiguring lesions. The greatest number of patients was in the age group of 61–80 years (47.2%) followed by 41–60 years (38.95%), 21–40 years (8.3%), and 81–100 years (5.6%), respectively . The youngest age of presentation in case of females was 29 years, while in males the corresponding age was 45 years. Correlation between gender and age group was not statistically significant (Fisher exact test value being 0.177), implying that these two variables are independent .

Out of all patients, 69.4% (25/36) hailed from rural areas. Majority of the patients were illiterate (80.6%) . A statistically significant association was seen between duration of disease and illiteracy (value = 6.95 and value = 0.01). This meant that illiterate patients present at a later stage of disease attributable to lack of awareness about disease entity. Farming was the main occupation among male patients (92.3%), while housekeeping was the major job among female patients (95.7%).

The duration of disease before seeking medical care ranged from 5 months to as long as 15 years, with mean duration being 4.7 years. The average duration of sun exposure was 6 hours/day in case of males and 2.91 hours/day in female patients. This difference in duration of sun exposure was statistically significant (-value being 6.71 and value = 0.000) . However, the females were intermittently exposed to high intensity sunlight

due to work in open kitchens and fields during sowing and harvesting seasons.

None of the patients had been taking photoprotective measures such as use of sunscreens and protective clothing. There was no history of treatment with PUVA or NBUVB in any of the study cases. All the patients were nonalcoholics and nonsmokers. No patient had features suggestive of genodermatoses associated with predilection for cutaneous malignancies like xeroderma pigmentosa, albinism, and so forth. Out of 36 patients, one (2.8%) had been previously treated for breast and endometrial carcinoma. Family history of cutaneous and systemic malignancies was not present in any of them. All the cases belonged to Fitzpatrick skin types III and IV (calculated via Fitzpatrick scoring scale).

The most common histopathological variant was nodular subtype (77.8%) with a significant proportion of tumors being pigmented (16.7%). Other subtypes included basosquamous (8.3%), micronodular (2.8%), morpheaform (2.8%), keratotic (2.8%), and adenoid (2.8%) BCC and BCC with adnexal differentiation (2.8%).

DISCUSSION

Basal cell carcinoma occurs worldwide. So far, BCC has been considered as the disease of the White¹⁴. Consequently, most of the studies have focused on White populations in Europe, USA, and Australia with scarcity of data from developing countries. Estimates of the incidence of BCC are imprecise since there is no cancer registry that collects data on BCC.

Although incidence rates of BCC vary significantly according to the ethnicity and geographic location, most studies show a rising trend in its incidence worldwide. This has been largely attributed to fair complexion and ozone layer depletion resulting in increased UV radiation reaching earth's surface. Similar increasing trend was noticed in our study as well. But factors other than the mentioned above need to be searched and verified as darker skin complexion in Indians should otherwise be protective against BCC. Moreover, ozone layer destruction is most evident over the temperate and polar regions, while India is a tropical country^[15, 16].

Basal cell carcinoma is rare in young populations.

An increased incidence has also been noticed in children and young adults^[17]. This finding highlights the need for early institution of UV protection and skin cancer screening in the pediatric and young adult population. However, there was no case below the age of 20 years in our study. Radiotherapy is another risk factor for the development of BCC in younger age group. Relative risk of BCC is more for children who have undergone radiation therapy for enlarged thymus^[18] or neoplasms such as medulloblastoma^[19].

BCCs are more common in males as reported in most studies worldwide, presumably due to greater occupational and recreational exposure to ultraviolet radiation (UVR). However, an unusual female preponderance was noticed in our study which is consistent with findings of another Indian series^[20]. Indian housewives especially rural women work in open kitchen during their household chores and work in the fields during sowing and harvesting seasons exposing them to intermittent, high intensity UVR. It might explain higher frequency of BCC in females in our study as intermittent rather than constant, cumulative UVR exposure is implicated in the pathogenesis of BCC^[21]. This female predilection may also be attributed to the changes in cultural practices like "veil" custom, structurally thinner skin with lower collagen density in the dermis when compared to men.

In our study, higher frequency among rural inhabitants was seen when compared to urban residents. This can be explained on the basis of more outdoor activities (as agriculture is the main occupation), changes in clothing preferences, illiteracy, and infrequent use of sunscreens. The rural patients regard initial lesions of BCC as a minor cosmetic problem with insignificant impact on health and seek medical advice only when lesions become symptomatic or disfiguring. So, late presentation to health facilities is equally contributory. A study done in Punjab regarding cancer found that tap water contains high content of arsenic, chromium, iron, and mercury, whereas ground water has abundance of arsenic, chromium, nickel, and iron. Even pesticides have been detected in the locally grown vegetables as well. Tseng et al. found a dose-dependent relation between arsenic levels in drinking water and the prevalence of skin cancers. Thus, exposure to harmful metals and pesticides may also add to the risk of skin

cancers, but further clinical and research studies are needed to confirm their role in the pathogenesis of BCC and to delineate underlying mechanisms. Occupations at risk of BCC that are highlighted in our study include agricultural workers and housekeepers.

Although most BCCs are slow-growing, relatively nonaggressive tumors, a minority have an aggressive behavior with local tissue destruction and, rarely, metastasis. Metastatic BCC has a reported incidence of only 0.0028–0.5%. Risk factors for development of metastatic BCC include large primary tumor (>2 cm), location in head and neck region, long standing lesion, multiple primary tumors and recurrences, prior radiation therapy, large tumor depth, invasion of perineural space and blood vessels, fair skin, male gender, and immunosuppression. One of our patients was detected with metastatic BCC.

Conclusion

This study highlights a paradoxically increasing trend of BCC with female predilection and higher percentage of pigmented lesions in Indians. This skin malignancy tends to be commoner in rural and agriculture based population. Major contributory risk factors include intermittent rather than constant UV exposure, cultural and lifestyle changes, cosmetic indifference, possible role of arsenic and pesticides, improved clinical awareness, and diagnostic facilities. The increasing cancer burden calls for the need of introduction of national screening program including mandatory annual skin examination by trained health professionals at the national level. Since early detection and treatment of lesions are crucial to decrease functional and cosmetic morbidity and costs, this study highlights the importance of improving awareness among general practitioners, public health workers, and general population. The clinical and epidemiological data collected in this study would serve as a reference for future research and may be helpful in the development of preventive and educational strategies.

Ethical Clearance- Taken from ethical committee of institution

Source of Funding- Self

Conflict of Interest – Nil

References

1. Malik V, Goh KS, Leong S, Tan A, Downey D, O'Donovan D. Risk and outcome analysis of 1832 consecutively excised basal cell carcinoma's in a tertiary referral plastic surgery unit. *Journal of plastic, reconstructive & aesthetic surgery*. 2010 Dec 1;63(12):2057-63.
2. Bradford PT. Skin cancer in skin of color. *Dermatology nursing/Dermatology Nurses' Association*. 2009 Jul;21(4):170. View at: Google Scholar
3. Deo SV, Hazarika S, Shukla NK, Kumar S, Kar M, Samaiya A. Surgical management of skin cancers: Experience from a regional cancer centre in North India. *Indian journal of cancer*. 2005 Jul 1;42(3):145. View at: Publisher Site | Google Scholar
4. HYMAN AB, BARSKY AJ. Basal cell epithelioma of the palm. *Archives of dermatology*. 1965 Nov 1;92(5):571-3. View at: Publisher Site | Google Scholar
5. Pearson G, King Jr LE, Boyd AS. Basal cell carcinoma of the lower extremities. *International journal of dermatology*. 1999 Nov;38(11):852-4. View at: Publisher Site | Google Scholar
6. Newman JC, Leffell DJ. Correlation of embryonic fusion planes with the anatomical distribution of basal cell carcinoma. *Dermatologic surgery*. 2007 Aug;33(8):957-65. View at: Publisher Site | Google Scholar
7. Markey AC, Lane EB, Macdonald DM, LEIGH IM. Keratin expression in basal cell carcinomas. *British Journal of Dermatology*. 1992 Feb;126(2):154-60. View at: Publisher Site | Google Scholar
8. LeSueur BW, Silvis NG, Hansen RC. Basal cell carcinoma in children: report of 3 cases. *Archives of dermatology*. 2000 Mar 1;136(3):370-2. View at: Google Scholar
9. Roudier-Pujol C, Auperin A, Nguyen T, Duvillard P, Benhamou E, Avril MF. Basal cell carcinoma in young adults: not more aggressive than in older patients. *Dermatology*. 1999;199(2):119-23. View at: Publisher Site | Google Scholar
10. Lim JL, Stern RS. High levels of ultraviolet B exposure increase the risk of non-melanoma skin cancer in psoralen and ultraviolet A-treated patients. *Journal of investigative dermatology*. 2005 Mar 1;124(3):505-13. View at: Publisher Site

| Google Scholar

11. M. R. Wehner, M. L. Shive, M.-M. Chren, J. Han, A. A. Qureshi, and E. Linos, "Indoor tanning and non-melanoma skin cancer: systematic review and meta-analysis," *British Medical Journal*, vol. 345, no. 7877, Article ID e5909, 2012.
12. Karagas MR. Occurrence of cutaneous basal cell and squamous cell malignancies among those with a prior history of skin cancer. *Journal of investigative dermatology*. 1994 Jun 1;102(6). View at: Google Scholar
13. Heal C, Buettner P, Browning S. Risk factors for wound infection after minor surgery in general practice. *Medical journal of Australia*. 2006 Sep;185(5):255-8. View at: Google Scholar
14. Lomas A, Leonardi-Bee J, Bath-Hextall F. A systematic review of worldwide incidence of nonmelanoma skin cancer. *British Journal of Dermatology*. 2012 May;166(5):1069-80. View at: Publisher Site | Google Scholar
15. "Myth: Ozone Depletion Occurs Only in Antarctica | Science | Ozone Layer Protection | US EPA," http://www.epa.gov/ozone/science/myths/glob_dep.html. View at: Google Scholar
16. Montzka SA, Reimannander S, Engel A, Kruger K, Sturges WT, Blake DR, Dorf MD, Fraser PJ, Froidevaux L, Jucks K, Kreher K. Ozone-Depleting Substances (ODSs) and Related Chemicals, Chapter 1 in *Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and Monitoring Project- Report No. 52*, 516 pp., World Meteorological Organization, Geneva, Switzerland, 2011. http://search.usa.gov/search?affiliate=esrl.noaa.gov_csd&query=assessment+of+ozone+depletion.
17. L. J. Christenson, T. A. Borrowman, C. M. Vachon et al., "Incidence of basal cell and squamous cell carcinomas in a population younger than 40 years," *The Journal of the American Medical Association*, vol. 294, no. 6, pp. 681–690, 2005.
18. N. Meibodi, M. Maleki, Z. Javidi, and Y. Nahidi, "Clinicopathological evaluation of radiation induced basal cell carcinoma," *Indian Journal of Dermatology*, vol. 53, no. 3, pp. 137–139, 2008. View at: Publisher Site | Google Scholar
19. M. Endo, K. Fujii, K. Sugita, K. Saito, Y. Kohno, and T. Miyashita, "Nationwide survey of nevoid basal cell carcinoma syndrome in Japan revealing the low frequency of basal cell carcinoma," *American Journal of Medical Genetics Part A*, vol. 158, no. 2, pp. 351–357, 2012. View at: Publisher Site | Google Scholar
20. Mahanthesh M, Rameshkumar K. Basal cell carcinoma: evaluation of clinical and histologic variables. *Indian Journal of Dermatology*. 2004 Jan 1;49(01):25. View at: Google Scholar
21. Laishram RS, Banerjee A, Punyabati P, Sharma LD. Pattern of skin malignancies in Manipur, India: A 5-year histopathological review. *Journal of Pakistan Association of Dermatology*. 2016 Dec 21;20(3):128-32. View at: Google Scholar