

# The Role of IL6 in Spontaneous Preterm Labour

Intisar Younis Mohammed Ibrahim

High Diploma in Gynecology and Obstetric /College of Medicin/Univircity of Baghdad M.B.Ch.B. D.G.O Mosul  
General Hospital

## Abstract

**Objective:** Preterm birth (PTB) and Preterm labour (PTL) (PTL is defined as ‘regular uterine contractions, which result in cervical changes before 37 Week of pregnancy. The main cause of neonatal morbidity and mortality with 5-18% of pregnancies was the preterm labour. Interleukin-6 (IL-6) amniotic fluid is an important cytokine used to identify intra-amniotic inflammation and those at risk to imminent premature delivery in patients with elevated AF IL-6. However, results of the traditional measurement method (ELISA) are generally unavailable in time in order to inform care. The aim of this study was to establish if intra-amniotic inflammation and/or infectious patients and those intended to spontaneous pre-term delivery among women with preterm labour and intact membranes can be identified through a point of care (POC) test or lateral-flow immunoassay for AF IL-6 concentration measurement.

**Methods:** In this study, one hundred and thirty-six singleton pregnant women who had symptoms of premature labor and amniocentesis were included. At the time of diagnosis, amniocentesis was carried out. In this study, the determination of AF (white blood cells, gram stain) was happened. By the results of AF culture, microbial invasion of the amniotic cavity (MIAC) was defined. EnzymeLinked Immunosorbent Assay (ELISA) and the lateral flow based immunoassay used to determine AF IL-6 concentrations. Define AF ELISA IL-6 as  $\geq 2600$  pg/ml as the primary result for intra- amniotic inflammation

**Results:** (1) The concentrations of AF IL-6 has been determined by a POC test with a specificity (91%), sensitivity (93%) and a positive likelihood ratio (10) where Intraamniotic inflammation identified with the threshold of 740 pg/ml; (2) The POC test and IL-6’s ELISA are also used to identify patients at risk of acute placenta inflammation, MIAC and impending spontaneous preterm delivery.

**Conclusion:** Intra-amniotic inflammation in women who present subsequently with spontaneous delivery before 34 week of gestation or those who will present with preterm and intact membranes can be identified by POC AF IL-6 test. It can be achieved within 20 minutes – this has important clinical implications and opens up opportunities for early diagnosis and treatment of intra-amniotic infection / inflammation.

**Keywords:** Interleukin-6, Preterm birth, Preterm labor, Neonatal mortality, Inflammation.

## Introduction

The leading cause of neonatal morbidity and mortality [2] can be considered preterm birth which affects on 5 – 18 % of pregnancies [1]. In the women who deliver preterm, one in four of them infected with an intraamniotic infection that is largely subclinical [3,4]. Inflammatory processes (that mediated Microbial associated preterm labour) involving production of

cytokines such as interleukin IL-6 [5,6], IL-1 [7], IL-10 [8], matrix degrading enzymes [9], chemokines [6], tumor necrosis factor-alpha (TNF- $\alpha$ ) [6,10] and other inflammatory-related proteins activate common parturition pathway [1]. Multiple studies showed that IL-6 amniotic fluid (AF) is superior to AF (glucose, Gram stain, white blood cell (WBC) counts) , or equal to proteomic markers for identification of the amniotic cavity intraamniotic and amniotic microbial invasions

(MIAC) [11]. In addition, the elevated concentrations of AF IL-6, even if the amniotic cavity absence the demonstrable microorganisms, is associated with an increase risk of neonatal outcomes and adverse pregnancy in the context of preterm prelabor rupture of the membranes (preterm PROM) [12], preterm labor [13] and a short cervix [14]. The concentrations of AF IL-6 therefore have both diagnostic and prognostic value.

The determination of AF IL-6 concentrations takes usually hours and results often cannot be made available on time in order to inform clinical decisions. In adult and neonatal sepsis [15] as well as in other inflammation-related conditions [16], there was widely used a POC (lateral flow-based immunoassay) test. Recently, these tests were applied in obstetrical applications. In the pilot study our group has found that AF IL-6 levels determined through POC tests correlated strongly with those measured through conventional immunosorbent (ELISA) (Spearman's  $\rho = 0.92$ ) [17]. In addition, our group has shown strong correlations. In addition, the POC IL-6 test results may identify preterm PROM patients intended for preterm delivery and/or acute histological chorioamnionitis [18].

In the present study determined by a POC test, we investigate whether AF IL-6 concentrations are able to identify patients (with intact membranes and preterm labour) who deliver spontaneously before term and/or have intraamniotic infection and or inflammations, relative to the concentrations performance determined by conventional ELISA.

## Materials and Methods

### Study populations

A retrospective cohort study used to identify patients with intact membranes with a diagnosis of spontaneous preterm labor. If the following criteria were met, the patients could be included: (1) microbiological studies conducted for 20 to 35 weeks of transabdominal amniocentesis (2); available AF for microbiological research performance (3) singleton gestation; and (4) neonatal results known. Patients who had a fetus with structural or chromosomal abnormality or they had

placenta previa, were excluded. Patients diagnosed with preterm labor associated with intact membranes, were told by their treating doctors about the potential value of microorganisms for AF identification. For research reasons outside of the clinical studies, women who agreed to undergo an amniocentesis were asked to donate AF. The doctor in charge of the further treatment of these patients. All patients received informed consent in writing and the use of clinical data and biological specimens were approved.

### Samples and biological analyzes

To the clinical laboratory for genital mycoplasma and (anaerobic, aerobic bacteria) and in a syringe with a capped sterile, AF was transported. The centrifuge of AF, that is not required for clinical assessment, was for 15 min at 4°C and stored at -75°C until analysis. The assessment was also carried out soon after gram stain, glucose, WBC count and AF were collected. To know if there is intraamniotic inflammations and/or infection, the concentrations of AF IL-6 needed and by ELISA was determined.

### Clinical definitions

Preterm labor was diagnosed in patients with cervical changes involving at least two regular uterine contractions every 10 minutes, related to gestational age from 20 to 36 6/7 weeks. Acute histologic chorioamnionitis was diagnosed with the above criteria [19]. Funisitis was diagnosed by the use of previously reported criterion for infiltration by neutrophils in walls of the umbilicus or Wharton jelly [20]. When the AF IL-6 concentration was  $\geq 2600$  pg/ml ( $\geq 2.6$  ng/ml), that determine by ELISA, Intra-amniotic inflammation has been diagnosed [21]. Based on the results of AF culture, MIAC was defined. The combination of intra-amniotic inflammation and MIAC was defined as intra-amniotic infection.

### IL-6 concentrations and analyzing it by AF samples

In the case of AF IL-6 concentrations (pg/ml), the Enzyme Linked Immunosorbent Assay and lateral immuno-assay POC tests were both determined. The immunoassays from R&D Systems for ELISA

have been used to determine AF IL-6 concentrations (Minneapolis, MN). A lateral flow-based immunoassay test POC was used to determine AF IL-6 concentrations (pg/ml). ELISA [6,11] and POC immunoassay details and performance have been described previously [17]. For IL-6 POC tests, the intra-assay variation coefficients and inter are 12.1% and 15.5% respectively.

### Study outcomes

The main results of this study are intra-amniotic inflammation and positive AF culture. While Spontaneous preterm (24 hour, 48 hour and 7days admission), acute inflammatory placental lesions (acute histologic chorioamnionitis and / or acute funisitis) with spontaneous preterm delivery (<28 week and <34 week of gestation) are considered the secondary results. The relation between acute chorioamnionitis histologically and amniotic fluid IL-6 concentrations were examined in 60 patients administered three days after amniocentesis. In this interval, a significant temporal relation between placental pathology and the amniocentesis results has been maintained.

### Statistical Analysis

In order to evaluate normality in arithmetic data distributions, the Kolmogorov–Smirnov test used. Kruskal–Wallis and Mann–Whitney U tests

were performed to compare the groups of arithmetic variables. Chi-square or Fisher's exact test was used for comparisons of the categorical variable. SPSS 19 (IBM Corp., Armonk, NY) and SAS 9.4 were used for statistical analysis (Cary, NC). Statistically significant was a p value <0.05.

## Results

### Characteristics of study populations

In this study, 136 women have been included with preterm labour with intact membranes. In Table 1, The clinical features are listed. The prevalence was nearly 16% (23/136) for MIAC and nearly 44% (60/136) for intra-amniotic inflammation. Spontaneous preterm deliveries were reported in most participants, nearly 22.7% (31/136) for < 28 weeks, 54.1% (74/136) for < 34 weeks and 64.2% (87/136) for < 37 weeks. In 24 hr, 48 hr, and 7 days, spontaneous delivery rates ranged between 33.7% (46/136), 43.5% (59/136) and 47.7% (65/136) respectively. In amniocentesis, the median gestational age was 30.8 weeks (27–32.4). 57.2% (31/54) of the 54 women with an acute histological chorioamnionitis and a large number of funisitis were diagnosed within 3 days after amniocentesis and placenta pathology reports [67.8% (21/31)].

**Table 1. ' Study populations' clinical features or characteristics**

Table 1. Clinical characteristics of the study population.	
Characteristic	Median (interquartile range) or percent (n = 136)
Maternal age (years)	24 (20–29)
Nulliparity	33.8% (46/136)
Prior preterm delivery	37.5% (51/136)
Gestational age at amniocentesis (weeks)	30.9 (27–32.4)
Amniotic fluid glucose (mg/dl)	24 (17–30.8)
Amniotic fluid white blood cell (cell/m <sup>3</sup> )	1.5 (0–13)
Microbial invasion of the amniotic cavity (%)	16.2% (22/136)
Intra-amniotic inflammation (ELISA IL-6 ≥ 2600 pg/ml) (%)	44.1% (60/136)
Gestational age at delivery (weeks)	33.3 (28.2–36.9)
Interval from amniocentesis to delivery (d)	8 (1–36.8)
Spontaneous deliver within one day after amniocentesis (%)	33.8% (46/136)
Spontaneous deliver within two days after amniocentesis (%)	43.4% (59/136)
Spontaneous deliver within seven days after amniocentesis (%)	47.8% (65/136)
Spontaneous delivery at <28 weeks of gestation (%)	22.8% (31/136)
Spontaneous delivery at <34 weeks of gestation (%)	54.4% (74/136)
Spontaneous delivery at <37 weeks of gestation (%)	64% (87/136)
Acute histologic chorioamnionitis (%)*	57.4% (31/54)
Acute funisitis (%)*	38.9% (21/54)
Acute inflammatory lesions of placenta (%)*	57.4% (31/54)

The data presented as % (n) or median. Inflammatory lesions of placenta acutely include acute chorioamnionitis histologically and acute funisitis which included only patients who had interval from amniocentesis to delivery < 4 d (n= 58).

and POC), inflammatory AF and placental lesion type or absence of acute inflammation in MIAC women are listed. *Ureaplasma urealyticum*, which was found in 18 percent (4/22) of these women, was the most often identified microorganism.


In Table 2, the microorganisms identified by culture AF, gestational aging at delivery, IL-6 levels (by ELISA

**Table 2. Inpatients with microbe invasion of the cavity using cultivation techniques, clinical characteristics, inflammatory amniotic fluid response, and acute placental inflammatory lesions.**

Table 2. Clinical characteristics, amniotic fluid inflammatory response and acute inflammatory placental lesions in patients with microbial invasion of the amniotic cavity using cultivation techniques.

No.	Organisms	GA at delivery (weeks)	AF glucose (mg/dl)	AF WBC (cell/mm <sup>3</sup> )	ELISA IL-6 (pg/ml)	Point of care IL-6 (pg/ml)	Acute histological chorioamnionitis	Acute funisitis
1.	<i>Prevotella</i> spp., <i>Enterococcus faecalis</i>	25 <sup>+1</sup>	10	1	52 637	3208	No	No
2.	<i>Mycoplasma hominis</i>	33	1	420	172 301	4613	N/A	N/A
3.	<i>Ureaplasma urealyticum</i>	28 <sup>+6</sup>	13	180	9433	10 000	Acute chorioamnionitis	No
4.	<i>Candida albicans</i>	26 <sup>+3</sup>	10	2160	201 261	4448	Necrotizing chorioamnionitis	No
5.	<i>Streptococcus agalactiae</i>	25	10	4	93 638	3575	Necrotizing chorioamnionitis	No
6.	<i>Candida albicans</i> , <i>Lactobacillus</i> spp.	33 <sup>+1</sup>	10	43	200 626	3554	Acute subchorionitis	No
7.	<i>Haemophilus influenza</i>	30 <sup>+6</sup>	10	40	92 063	6467	Necrotizing chorioamnionitis	Necrotizing funisitis
8.	<i>Fusobacterium</i> spp., Gram-negative bacilli	21 <sup>+6</sup>	19	1564	317 655	6228	Subchorionic microabscesses	Necrotizing funisitis
9.	Gram-negative bacilli	21 <sup>+1</sup>	20	66	242 699	5934	Subacute chorioamnionitis	Umbilical arteritis
10.	<i>Streptococcus agalactiae</i>	25 <sup>+2</sup>	19	5	248 889	8208	Acute chorioamnionitis	Umbilical arteritis
11.	Gram-positive cocci	22 <sup>+5</sup>	10	125	470 626	5540	Necrotizing chorioamnionitis	Umbilical arteritis
12.	<i>Fusobacterium</i> spp., Gram-negative bacilli	28 <sup>+1</sup>	1	22	301 426	3996	Acute chorioamnionitis	Umbilical arteritis
13.	<i>Ureaplasma urealyticum</i>	33	10	500	85 962	4628	Acute chorioamnionitis	Umbilical phlebitis/chorionic vasculitis
14.	<i>Streptococcus anginosus</i> , <i>Streptococcus mitis</i>	22 <sup>+6</sup>	10	10	73 254	7246	Acute chorioamnionitis	Umbilical phlebitis/chorionic vasculitis
15.	<i>Bacteroides</i> spp., <i>Mobiluncus</i> spp., <i>Clostridium sporogenes</i>	22 <sup>+4</sup>	10	295	517 846	4748	Necrotizing chorioamnionitis	Umbilical phlebitis/chorionic vasculitis
16.	<i>Ureaplasma urealyticum</i>	39 <sup>+2</sup>	19	2	1274	155	No	No
17.	Gram-negative bacilli	26 <sup>+5</sup>	N/A	610	1779	10 000	N/A	N/A
18.	<i>Streptococcus</i> spp., <i>Gemella morbillorum</i>	31 <sup>+6</sup>	10	1920	741	6796	Necrotizing chorioamnionitis	Umbilical arteritis
19.	<i>Candida albicans</i>	32 <sup>+4</sup>	10	1292	96 334	4252	Acute chorioamnionitis	Necrotizing funisitis
20.	<i>Staphylococcus capitis</i>	28 <sup>+6</sup>	20	24	360 503	4374	Subacute chorioamnionitis	Umbilical arteritis
21.	<i>Mobiluncus</i> spp.	32	10	570	76 888	8144	Acute chorioamnionitis	Umbilical phlebitis/chorionic vasculitis
22.	<i>Ureaplasma urealyticum</i>	34 <sup>+3</sup>	24	0	244	60	Acute subchorionitis	No

N/A: results were not available; WBC, white blood cell count; AF, amniotic fluid: acute subchorionitis/chorionitis = acute histologic chorioamnionitis stage 1; acute chorioamnionitis = acute histologic chorioamnionitis stage 2; necrotizing chorioamnionitis and subacute chorioamnionitis = acute histologic chorioamnionitis stage 3: subchorionic microabscesses = severe acute histologic chorioamnionitis; umbilical phlebitis/chorionic vasculitis = acute funisitis stage 1; umbilical arteritis = acute funisitis stage two; necrotizing funisitis = stage three acute funisitis.



**Figure 1. Characteristic curve of the operating receiver describes the performance of an amnioticFluid Interleukin-6 test in intra-amniotic inflammation identification ( the area under curve for amniotic fluid IL-6 point =0.93; 96% confidence interval: 0.91–0.98, P<0.002).**

**The diagnosis of an AF IL-6 POC test to identify the inflammation of the intraamniotic** A threshold of = 745 pg/ml was selected for the POC test after inspecting a recipient operationalcharacteristic curve to identify an intraamniotic inflammation [zone of curve = 0.94 (0.90–0.99)]. (Figure 1). The POC test was given a sensitivity of 93% and a specialty of 91% in Table 3 lists.

The diagnostic performance of an AF IL-6 POC test for the identification of MIAC and acute inflammatory lesions of placenta

Table 3 shows that the POC test performance was equivalent to conventional ELISA when identifying MIAC and placental lesion patients in line with

acute inflammation. 18.2% (4/22) of MIAC patients showed negative results for ELISA IL-6 (Table 2). In two of these four patients, however, the POC AF IL-6 was raised. One of those patients with acute histological chorioamnionitis and funisitis implies a real intraamniotic infection. A placental pathology reportwas not provided to the other patient. Note, the potential for a contamination is suggested by one of two patients who had AF IL-6 negatives in both trials (POC and ELISA) delivered at the term of the trial and had not placental lesions consistent with acute inflammation.

The identification of preterm delivery The AF IL-6 POC test

In **Table 3**, the tests for women spontaneously deliveries are presented with Point of care test and Enzyme Linked Immunosorbent Assay AF IL-6. Table 2. Both tests had positive probability equivalents in identifying patients who had spontaneous preterm delivery within one of the days of amniocentesis or patients who spontaneously delivered after a gestation of less than 28 weeks. Each one had marginally a greater sensitivity and specificity than ELISA for the POC test in determining women who would spontaneously deliver within two or seven days of amniocentesis. In comparison with ELISA's results the sensitivity for spontaneous preterm delivery was slightly higher at less than 34 weeks of gestation, while the specificity was slightly less. However, confidence intervals in the assessment of the diagnostic performance of the POC test have overlapped the ELISA test, statistic equivalent performance in the assessment of the risk of spontaneous premature delivery.

### Discussion

The main findings from the study: (1) The POC-determined AF-IL-6 concentrations are highly sensitive (93%) and are specific (91%) to identify intra-amniotic inflammation with a 745 pg/mL threshold and (2) IL-6 determination by POC test and ELISA was performed similarly in patients identifications with acute inflammatory lesions of placenta, MIAC and the patient with intact membrane and preterm labour at risk of spontaneous preterm delivery.

**AF IL-6 POC test for intra-amniotic inflammation identification and imminent preterm delivery** Preterm labor and other adverse effects are at greater risk of intraamniotic inflammation, as demonstrated by compelling evidence without identifiable microorganisms<sup>[13]</sup>. In patients with preterm labor and intact membranes, sterile intra-amniotic inflammation, an inflammatory process in which neither culture nor molecular methodology can be observed, has previously been demonstrated to be more prevalent than intra-amniotic, microbial inflammation.<sup>[22]</sup> preterm PROM and asymptomatic sonographic short cervix<sup>[14]</sup>. We have also shown that sterile inflammation intraamniotic is connected to unhealthy pregnancy; hence intraamniotic

inflammation is important<sup>[22, 14]</sup>.

In this study, we have shown that the point of care AF IL-6 test is highly sensitive and specific to identify spontaneous premature delivery and intraamniotic inflammations. The results of the POC test are compared with AF IL-6 concentrations determined by ELISA in the identification of infection-inflammation-related obstetric outcomes, but they could be determined within 20 minutes. Therefore, contrary to conventional ELISA, the POC AF IL-6 results can be available on time for clinical decisions, similarly to an MMP 8 rapid matrix test which has demonstrated that patients with preterm labor and intact membranes with >80 percent susceptibility and >90 percent specificity have an intraamniotic infection or inflammation<sup>[23]</sup>. In addition, the MMP-8 test has been shown to be useful in identifying intra-amniotic inflammation in preterm PROM, MIAC in preterm delivery risk patients, and funisitis in preterm delivery patients.

Interestingly, six out of seven patients with POC and ELISA negative results had spontaneous early (<34 weeks) preterm delivery within a period of two days after amniocentesis. This suggests that the POC test provides additional risk information beyond the standard ELISA tests. It should also be noted that two out of four POC negative and ELISA positive which at term delivered, one of whom did not have acute inflammation with placental lesions and the result was positive (2609 pg/ml). In the previous study, we showed that the concentrations of AF IL-6 that determined by POC test were 30 percent less on average than those established by conventional ELISA. Therefore, it is no surprise that the POC selected the lower AF IL-6 cut-off for intra amniotic

inflammation (THI) patients (THI 745 pg/ml) in this study). Kacerovsky and others suggested a higher reduction (1000 pg/ml) to detect MIAC in a study that used the same test for AF IL 6 levels among preterm MIAC women or the combination of MIAC with acute histological chorioamnionitis. Other POC test researchers reported a high negativity of IL-6 detection predictive value in vaginal fluid in women with preterm PROM, comparable to that found in our study (97.6 percent versus

94.4 percent ) [23]. However, our study showed more positive predictive value than the concentration of IL-6 vaginal fluid POC test (88.9 percent versus 50 percent ). Using POC IL-6 in vaginal fluid for pregnancy results has been reported by Vousden et al. in asymptomatic high risk patients with preterm birth [18]. A cut off of 56 pg/ml had 81 percent sensitivity and 65 percent specificity for the vaginal fluid IL-6 concentration for identifying people who were given under 28 weeks of gestation [18]. The diagnostic performances in this study are somewhat lower than POC AF IL-6. In line with the risk/benefit ratios for specific actions, the optimal cutoff value is to be determined.

### Strengths and Limitations

The study strengths include: - (1) Test POC for the purpose of informing treatment was not used and (2) including a group of patients with preterm labour with intact membrane instead of patient with preterm prelabour rupture of membranes who have high prevalence of infection and/or inflammation. One constraint is the use of cultivation techniques to identify amniotic cavity microorganisms, therefore, it may not be possible to detect non-cultural bacteria that could have been identified using molecular microbiological techniques.

### Conclusions

Intra-amniotic inflammation can be identified by a POC AF IL-6 test as determined by ELISA in women with premature labour and intact membranes and also by equivalently identifying those who deliver spontaneously before the term subsequent to this test. Further studies are necessary to determine whether the results of POC AF IL-6 inform sufficient therapeutic decisions to improve pregnancy results in patients of this type.

**Conflict of Interest:** There is no conflict of interest among the authors.

**Funding:** Self

**Ethical Clearance:** This study is ethically approved by the Institutional ethical Committee.

### References

1. Romero R, Dey SK, Fisher SJ. Preterm labor: one syndrome, many causes. *Science*. 2014 Aug 15;345(6198):760-5.
2. Landmann E, Misselwitz B, Steiss JO, Gortner L. Mortality and morbidity of neonates born at < 26 weeks of gestation (1998–2003). A population-based study. *Journal of perinatal medicine*. 2008 Mar 1;36(2):168-74.
3. Bobitt JR, Ledger WJ. Unrecognized amnionitis and prematurity: a preliminary report. *The Journal of reproductive medicine*. 1977 Jul 1;19(1):8-12.
4. Romero R, Espinoza J, Gonçalves LF, Kusanovic JP, Friel L, Hassan S. The role of inflammation and infection in preterm birth. In *Seminars in reproductive medicine* 2007 Jan (Vol. 25, No. 01, pp. 021-039). Copyright© 2007 by Thieme Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA..
5. Romero R, Avila C, Santhanam U, Sehgal PB. Amniotic fluid interleukin 6 in preterm labor. Association with infection. *The Journal of clinical investigation*. 1990 May 1;85(5):1392-400.
6. Chaemsaihong P, Romero R, Korzeniewski SJ, Martinez-Varea A, Dong Z, Yoon BH, Hassan SS, Chaiworapongsa T, Yeo L. A rapid interleukin-6 bedside test for the identification of intra-amniotic inflammation in preterm labor with intact membranes. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2016 Feb 1;29(3):349-59.
7. Mackler AM, Iezza G, Akin MR, McMillan P, Yellon SM. Macrophage trafficking in the uterus and cervix precedes parturition in the mouse. *Biology of reproduction*. 1999 Oct 1;61(4):879-83.
8. Gotsch F, Romero R, Kusanovic JP, Erez O, Espinoza J, Kim CJ, Vaisbuch E, Than NG, Mazaki-Tovi S, Chaiworapongsa T, Mazor M. The anti-inflammatory limb of the immune response in preterm labor, intra-amniotic infection/inflammation, and spontaneous parturition at term: a role for interleukin-10. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2008 Jan 1;21(8):529-47.
9. Athayde N, Romero R, Gomez R, Maymon E, Pacora P, Mazor M, Yoon BH, Fortunato S, Menon

- R, Ghezzi F, Edwin SS. Matrix metalloproteinases-9 in preterm and term human parturition. *The Journal of Maternal-Fetal Medicine*. 1999 Sep;8(5):213-9.
10. Armstrong-Wells J, Donnelly M, Post MD, Manco-Johnson MJ, Winn VD, Sébire G. Inflammatory predictors of neurologic disability after preterm premature rupture of membranes. *American journal of obstetrics and gynecology*. 2015 Feb 1;212(2):212-e1.
  11. Romero R, Yoon BH, Kenney JS, Gomez R, Allison AC, Sehgal PB. Amniotic fluid interleukin-6 determinations are of diagnostic and prognostic value in preterm labor. *American journal of reproductive immunology*. 1993 Sep 10;30(2-3):167-83.
  12. Yoon BH, Romero R, Moon JB, Shim SS, Kim M, Kim G, Jun JK. Clinical significance of intra-amniotic inflammation in patients with preterm labor and intact membranes. *American journal of obstetrics and gynecology*. 2001 Nov 1;185(5):1130-6.
  13. Romero R, Miranda J, Chaiworapongsa T, Chaemsaihong P, Gotsch F, Dong Z, Ahmed AI, Yoon BH, Hassan SS, Kim CJ, Korzeniewski SJ. Sterile intra-amniotic inflammation in asymptomatic patients with a sonographic short cervix: prevalence and clinical significance. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2015 Jul 24;28(11):1343-59.
  14. Meem M, Modak JK, Mortuza R, Morshed M, Islam MS, Saha SK. Biomarkers for diagnosis of neonatal infections: A systematic analysis of their potential as a point-of-care diagnostics. *Journal of global health*. 2011 Dec;1(2):201.
  15. Dengler J, Schefold JC, Graetz D, Meisel C, Splettstößer G, Volk HD, Schlosser HG. Point-of-care testing for interleukin-6 in cerebro spinal fluid (CSF) after subarachnoid haemorrhage. *Medical Science Monitor*. 2008 Dec 1;14(12):BR265-8.
  16. Chaemsaihong P, Romero R, Korzeniewski SJ, Dong Z, Yeo L, Hassan SS, Kim YM, Yoon BH, Chaiworapongsa T. A point of care test for the determination of amniotic fluid interleukin-6 and the chemokine CXCL-10/IP-10. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2015 Sep 2;28(13):1510-9.
  17. Vousden N, Chandiramani M, Seed P, Shennan A. Interleukin-6 bedside testing in women at high risk of preterm birth. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2011 Oct 1;24(10):1301-4.
  18. Redline RW, Heller D, Keating S, Kingdom J. Placental diagnostic criteria and clinical correlation—a workshop report. *Placenta*. 2005 Apr 1;26:S114-7.
  19. Pacora P, Chaiworapongsa T, Maymon E, Kim YM, Gomez R, Yoon BH, Ghezzi F, Berry SM, Qureshi F, Jacques SM, Kim JC. Funisitis and chorionic vasculitis: the histological counterpart of the fetal inflammatory response syndrome. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2002 Jan 1;11(1):18-25.
  20. Romero R, Chaiworapongsa T, Alpay Savasan Z, Xu Y, Hussein Y, Dong Z, Kusanovic JP, Kim CJ, Hassan SS. Damage-associated molecular patterns (DAMPs) in preterm labor with intact membranes and preterm PROM: a study of the alarmin HMGB1. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2011 Dec 1;24(12):1444-55.
  21. Romero R, Miranda J, Chaiworapongsa T, Korzeniewski SJ, Chaemsaihong P, Gotsch F, Dong Z, Ahmed AI, Yoon BH, Hassan SS, Kim CJ. Prevalence and clinical significance of sterile intra-amniotic inflammation in patients with preterm labor and intact membranes. *American journal of reproductive immunology*. 2014 Nov;72(5):458-74.
  22. Berthiaume M, Rousseau É, Rola-Pleszczynski M, Pasquier JC. Rapid evaluation of the absence of inflammation after rupture of membranes. *The Journal of Maternal-Fetal & Neonatal Medicine*. 2014 Jun 1;27(9):865-9.
  23. Nien JK, Yoon BH, Espinoza J, Kusanovic JP, Erez O, Soto E, Richani K, Gomez R, Hassan S, Mazor M, Edwin S. A rapid MMP-8 bedside test for the detection of intra-amniotic inflammation identifies patients at risk for imminent preterm delivery. *American journal of obstetrics and gynecology*. 2006 Oct 1;195(4):1025-30.