

Effectiveness of Planned Teaching Program on Knowledge Regarding the Management of Hospital acquired Infections in Children.

Shivateerthayya Hiremath¹, Prasad Patil², Madusudhan K.G³, Princely B Olickal⁴

¹Professor, Department of Medical-Surgical Nursing, V.M.P. College of Nursing & Medical Research Institute, Akluj, Solapur, Maharashtra-413118, India.

²Vice Principal, Department of Child Health Nursing, V.M.P. College of Nursing & Medical Research Institute, Akluj, Solapur, Maharashtra-413118, India.

³Associate Professor, Department of Community Health Nursing, MIT Nursing College, Aurangabad, Maharashtra-431001, India.

⁴Associate Professor, Department of Child Health Nursing, MIT Nursing College, Aurangabad, Maharashtra-431001, India.

How to cite this article: Shivateerthayya Hiremath, Prasad Patil, Madusudhan K.G, Princely B Olickal et al Effectiveness of Planned Teaching Program on Knowledge Regarding the Management of Hospital acquired Infections in Children. Volume 13 | Issue 4 | October-December 2022

Abstract

In the present study, data were collected on knowledge regarding management of Hospital Acquired infections in children. The objective of the study is to identify the knowledge of B.Sc Nursing students regarding the management of Hospital acquired infections in children and to determine the association between post-test knowledge scores with selected demographic variables. The research method adopted for this study is the evaluative approach. The research design adopted for this study was a pre-experimental design with one group pretest and post-test design. A simple random technique was used to select the sample for the study. The sample consists of 60 B.Sc Nursing students of Nursing institutions. The instrument used for the data collection is a structured questionnaire. Collected data were analyzed by using descriptive and inferential statistics. The results of the study showed that the pretest overall knowledge score regarding various aspects of management of Hospital Acquired infections in children was a mean percentage of 46.86% and SD was 3.51. During the post-test overall knowledge score was 80.53% and SD was 2.98. Hence the difference between pretest and post-test over a knowledge score was 33.67. The results show that the difference between the pretest and post-test knowledge scores for the management of Hospital Acquired infections in children is statistically significant and this difference is due to planned teaching program in the management of Hospital Acquired infections in children. The conclusion was drawn on the basis of the following findings of the study. This study shows that there was a significant improvement in the knowledge of management of infections in children among B.Sc Nursing students as evidenced by pre-test and post-test knowledge scores.

Keywords: Asses, Knowledge, PTP, Hospital acquired infections

Introduction

A hospital-acquired infection is usually one that first appears three days after a patient is admitted to a hospital or other health care facility. Infections acquired in a hospital are also called Hospital acquired infections. About 5-10% of patients admitted to hospitals in the United States develop a Hospital Acquired infections. About 90,000 of these patients

die as a result of their infections. Hospital-acquired infections usually are related to a procedure or treatment used to diagnose or treat the patient's illness or injury. About 25% of these infections can be prevented by healthcare workers taking proper precautions when caring for patients¹.

Hospital-acquired infections can be caused by bacteria, viruses, fungi, or parasites. These

Corresponding Author:

Shivateerthayya Hiremath

Professor, Department of Medical-Surgical Nursing, V.M.P. College of Nursing & Medical Research Institute, Akluj, Solapur, Maharashtra-413118, India.

E-mail: Shivateerth@gmail.com

microorganisms may already be present in the patient's body or may come from the environment, contaminated hospital equipment, health care workers, or other patients. Depending on the causal agents involved, an infection may start in any part of the body. A localized infection is limited to a specific part of the body and has local symptoms. For example, if a surgical wound in the abdomen becomes infected, the area of the wound becomes red, hot, and painful. A generalized infection is one that enters the bloodstream and causes general systemic symptoms such as fever, chills, low blood pressure, or mental confusion. Hospital-acquired infections may develop from surgical procedures, catheters placed in the urinary tract or blood vessels, or material from the nose or mouth that is inhaled into the lungs. The most common types of hospital-acquired infections are urinary tract infections (U.T.I.s), pneumonia, and surgical wound infections².

All hospitalized patients are susceptible to contracting a Hospital Acquired infections. Some patients are at greater risk than others young children, the elderly, and persons with compromised immune systems are more likely to get an infection. Other risk factors for getting a hospital-acquired infection are a long hospital stay, the use of indwelling catheters, failure of healthcare workers to wash their hands, and overuse of antibiotics³. Urinary tract infection (U.T.I.) is the most common type of hospital-acquired infection. Most hospital-acquired U.T.I.s happen after urinary catheterization. Catheterization is the placement of a catheter through the urethra into the urinary bladder. This procedure is done to empty urine from the bladder, relieve pressure in the bladder, measure urine in the bladder, put medicine into the bladder, or for other medical reasons. The healthy urinary bladder is sterile, which means it doesn't have any bacteria or other microorganisms in it. There may be bacteria in or around the urethra, but they usually cannot enter the bladder. A catheter can pick up bacteria from the urethra and allow them into the bladder, causing an infection to start⁴.

Bacteria from the intestinal tract are the most common type to cause U.T.I.s. Patients with poorly functioning immune systems or who are taking antibiotics are also at risk for infection by a fungus called *Candida*. Pneumonia is the second most common type of hospital-acquired infection. Bacteria and other microorganisms are easily brought into the throat by respiratory procedures commonly

done in the hospital. The microorganisms come from contaminated equipment or the hands of health care workers. Some of these procedures are respiratory intubation, suctioning of material from the throat and mouth, and mechanical ventilation. The introduced microorganisms quickly colonize the throat area. This means that they grow and form a colony but do not yet cause an infection. Once the throat is colonized, it is easy for a patient to inhale the microorganisms into the lungs. Patients who cannot cough or gag very well are most likely to inhale colonized microorganisms into their lungs. Some respiratory procedures can keep patients from gagging or coughing. Patients who are sedated or who lose consciousness may also be unable to cough or gag. The inhaled microorganisms grow in the lungs and cause an infection that can lead to pneumonia⁵.

WHO reported that every minute a mother dies from complications in pregnancy and childbirth which means 1400 mothers die every day. More than half a million mothers die every year. The birth of a baby is a momentous occasion in a couple's life. The aim of modern MANAGEMENT of labor should be to ensure optimum conditions for the mother and the fetus during and after delivery as well as emotional satisfaction for all involved. The place of delivery plays an important role in child survival and safe motherhood. A child born with an unhygienic condition is more to get the infection. A properly assisted delivery with skilled personnel and following aseptic precaution is highly advantageous to both mother and fetus during delivery⁶. Health can neither be demanded nor given, it can neither be bought nor sold, but the circumstances and services that are prerequisites to health can centrally be demanded and received as right. A protective environment in the hospital unit is essential as a prerequisite, particularly when considering the services provided in the labor room provision for a safe and protective environment is a priority need⁷.

Healthcare-associated infections are defined as infections not present and without evidence of incubation at the time of admission to a healthcare setting. To better reflect the diverse healthcare settings currently available to patients, the term healthcare-associated infections replaced old ones such as Nosocomial, hospital-acquired, or hospital-onset infections. Within hours after admission, a patient's flora begins to acquire the surrounding bacterial pool characteristics. Most infections that become

clinically evident after 48 hours of hospitalization are considered hospital-acquired. Infections that occur after the patient is discharged from the hospital can be considered healthcare-associated if the organisms were acquired during the hospital stay⁸.

Need for the Study

Today's Nursing students are tomorrow's staff nurses who can contribute themselves more in the field of treatment. But today, this nursing curriculum doesn't give much importance to infection control measures. Thus, students have less exposure to that field makes them vulnerable to knowledge. Educating these students and creating awareness in help them to learn more about the control of Nosocomial infections in children, and they will be able to control the infection in the pediatric hospital. In 2010 M.M.R. was reduced to less than 100. The suggestion was three antenatal check-ups, three postnatal check-ups, and clean, safe delivery given by the national health and family welfare department. Service to make motherhood safe includes care by skilled health personnel, aseptic delivery before, during, and after childbirth, health education for women, their families, and decision-makers; most maternal deaths and pregnancy complications can be prevented if efficient care is given during child care⁹.

Surgical procedures increase a patient's risk of getting an infection in the hospital. Surgery directly invades the patient's body, giving bacteria a way into normally sterile parts of the body. An infection can be acquired from contaminated surgical equipment or from healthcare workers. Following surgery, the surgical wound can become infected. Other wounds from trauma, burns, and ulcers may also become infected¹⁰. Many hospitalized patients need a steady supply of medications or nutrients delivered to their bloodstream. An intravenous (IV) catheter is placed in a vein, and the medication or other substance is infused into the vein. Bacteria transmitted from the surroundings, contaminated equipment, or healthcare workers' hands can invade the site where the catheter is inserted. A local infection may develop in the skin around the catheter. The bacteria also can enter the blood through the vein and cause a generalized infection. A localized infection causes swelling, redness, and tenderness at the site of infection¹¹.

Nursing personnel working in the pediatric ward should be knowledgeable and skillful in the management of Hospital Acquired infections. If the

nursing personnel fails to adopt the infection control technique, it may lead to septicemia and children's deaths. In order to prevent I.M.R., the nursing students (at the base level) should be knowledgeable in providing infection-free nursing care. The nurse concerned with the care and management of infection occurring through all the sources holds a responsible involvement position. They have an important role in the management of infection in a pediatric hospital. The nursing student has to be knowledgeable regarding preventive measures in controlling Hospital acquired infections¹².

Hospitals and other healthcare facilities have developed extensive infection control programs to prevent Hospital acquired infections. These programs focus on identifying high-risk procedures and other possible sources of infection. High-risk procedures such as urinary catheterization should be performed only when necessary, and catheters should be left in for as little time as possible. Medical instruments and equipment must be properly sterilized to ensure they are not contaminated. Frequent handwashing by healthcare workers and visitors is necessary to avoid passing infectious microorganisms to hospitalized patients. In 2003, the Joint Commission on Accreditation of Healthcare Organizations (JCAHO) announced it would make the management of Hospital acquired infections a major goal in 2004 and the coming years. JCAHO, the body that inspects hospitals for quality and accredits them accordingly, issued an alert stating that hospital-acquired infections are seriously underreported¹³.

The Study on the Efficacy of Hospital Acquired infections Control Project (SENIC) from 2005 showed Nosocomial rates could be reduced by 32% if infection surveillance were coupled with appropriate infection control programs. In 2005, Continued surveillance and sound infection control programs led to decreased healthcare-associated infections and better prioritization of resources and efforts to improve medical care. Healthcare-associated infections are of important wide-ranging concern in the medical field. They can be localized or systemic, involve any body system, and be associated with medical devices or blood product transfusions¹⁴.

The investigator observed that many times nursing students have adequate knowledge of aseptic techniques in a pediatric hospital. Hand washing, the single most measure in management and spread of infection from one person to another, is not being

done often. Another important area of concern is that nursing students should be careful in doing the vaginal examination, conducting delivery, aseptic precaution is very important to prevent the introduction of infection to the mother. The recognition of ignorance is the beginning of wisdom. To understand the degree of knowledge to help in future improvement thus, the investigator felt the need to take up the study such as it could assess the knowledge among B.Sc.Nursing students in selected Nursing institutions.

Objectives

The main objectives of the present study are mentioned here

Objectives of the Study

- To assess the pretest knowledge regarding the management of Hospital acquired infections in children.
- To educate the students with a planned teaching program on the management of Hospital acquired infections in children among B.Sc.Nursing students.
- To assess the effectiveness of a planned teaching program on the management of Hospital Acquired infections among B.Sc. Nursing students in terms of knowledge.
- To determine the association between post-test knowledge scores with selected demographic variables.

Operational Definitions

Assessment: It is the organized, systematic, and continuous process of collecting data regarding management of Hospital acquired infections in children.

Effectiveness: In this study, it refers to the extent to which the planned teaching program on the management of Hospital acquired infections in children has achieved the desired effect in improving the knowledge of B.Sc.Nursing students as evident from the gain in knowledge score.

Knowledge: It refers to the level of awareness among selected subjects regarding the management of Hospital acquired infections in children.

Hospital Acquired Infections: An infection acquired in the hospital by a patient who was admitted for a reason other than that infection.

B.Sc. Nursing Students: It refers to those students who are studying nursing course in selected Nursing institutions.

Variables

Dependent variable: In this study knowledge of B.Sc. Nursing students on management of Hospital acquired infections in children will be the dependent variable.

Attribute variable: In this study Age, gender, Religion, Type of family Occupation of the father and sources of information on the management of Hospital acquired infections in children are attribute variables.

Assumptions

- Students of B.Sc.Nursing may have inadequate knowledge on management of Hospital acquired infections in children.
- B.Sc. Nursing students may utilize the planned teaching program on management of Hospital acquired infections in children.
- A carefully prepared Planned Teaching Program may enhance the knowledge of B.Sc. Nursing students on management of Hospital acquired infections in children.

Hypothesis

H₁: There will be significant difference in the pretest and post knowledge score of B.Sc.Nursing students regarding management of Hospital acquired infections in children.

H₂: There will be a significant association between the post-test score with the selected demographic variables.

Limitations

- The study is limited to only the selected school of Nursing at Akluj.
- Students who are available at the time of data collection.
- The sample size is limited to 60 B.Sc.Nursing students from selected Nursing institutions at Akluj.

Research Methodology

Research Approach:

The researcher found that the **evaluative approach** is best suited and adopted for the present study.

Research Design

The present study's pre-experimental one group pretest and post-test design was selected.

Setting of the Study

The study was conducted at the school of Nursing setup.

Population

The population of the present study comprises selected Nursing institutions at Akluj. The accessible populations are those available at the time of conducting the study.

Sample

In the present study, 60 students of B.Sc.Nursing, studying in various institutions of Nursing, were selected.

Sampling Technique:

The sampling technique adopted to select the study samples was **simple random technique**.

Sampling Criteria

Inclusion Criteria

- Students who are willing to participate in the study.
- Both male and female students are included.
- B.Sc.Nursing. students studying at the selected Nursing institutions at Akluj.

Exclusion Criteria

- Students who are in internship.
- Students who are not available at the time of the study.

Method of Data Collection

Data Collection Instruments.

The present study is aimed at assessing the knowledge of students on the management of Hospital acquired infections in children. Thus, a structured knowledge questionnaire to assess the knowledge was prepared and used for data collection.

Development of Tool

A structured knowledge questionnaire was developed for assessing the knowledge on the management of Hospital acquired infections in children.

The tools were prepared on the basis of the objectives of the study.

Description of the Tool

Structured Knowledge Questionnaire

The structured knowledge questionnaire consists of two sections

Part - I

This section includes a demographic background of the students i.e. Age, Gender, Permanent area of residence, Previous academic performance, Religion, and Sources of information on infection control.

Part -II

This section is the second part of self-structured questionnaire, which consists of the following headings

Section - A: Consists of questions assessing knowledge about management of Hospital acquired infections in children's.

There are total 30 Multiple Choice Questions in section A, Item number 1-8 inquire general information on Hospital acquired infections in children, and 8-13 are related to measures used in the management of Hospital acquired infections in children, 14-25 sepsis and barrier techniques.

There are total of 30 questions in section B, of which are positively stated

Scoring

The 30 questions in section- A are Multiple-choice questions and for the correct option, the score is 1 and other options 0.

The 30 questions in Section- B are closed-ended dichotomous questions.

In positively stated items the score for Yes is 1 and for No is 0. In negatively stated items the score for Yes is 0 and for No is 1.

Data Collection Procedure

Prior to data collection permission was obtained from the concerned authorities. Further, the investigator obtained consent from the subjects. Confidentiality was maintained during data collection.

The data collection procedure was carried out for a period of six weeks. After obtaining permission from the authorities of a Nursing institutions Akluj. The

data collection was carried . After obtaining formal permission from the school of Nursing and from the participants, data were collected from 10 nursing students selected by a simple random sampling technique.

The investigator selected the Nursing students from VMP Nursing college at Akluj and administered the structured questionnaire to each of the B.Sc Nursing students.

It took about 1 hour to collect the data. The responses were recorded in the space provided in the questionnaire itself followed by a structured teaching program for the samples.

Results

Analysis and interpretation of the information collected through a structured questionnaire from 60 I year B.Sc.Nursing students of selected Nursing institutions at Akluj. The present study was designed to assess the effectiveness of Planned teaching program on knowledge management of Hospital acquired infections in children among collected data were coded, tabulated organized, analyzed, and interpreted using descriptive and inferential statistics.

Organizations of findings

The data collected from the B.Sc.Nursing.students were organized, analyzed and presented under the following headings.

- Section I: Description of sample characteristics
- Section II: Assessment of pre-existing knowledge.
- Section III: Evaluating the effectiveness of Planned teaching programme.
- Section IV: Association of Post-test knowledge with selected demographic variables.

Section-I

Demographic Characteristics of the Samples

It depicts that the majority of 47% were between the age group of 21-22yrs. Similarly, the remaining 26% of students were between the age group of 23-24 years, 18% of students were between the age group of 19-20yrs, and 8% were 24yrs and above. Gender shows that most nursing students were female, 58% and 42% were males. The majority of students, 53% were Christian, 23% students were Hindus, 17% were Muslims, and the remaining 7% of students

belonged to some other religion. Family depicts that the majority of 3rd year G.N.M.students were 47% from a joint family, 28% from a nuclear family, and the remaining 25% were from extended family. Works depict that the majority of students' fathers were, 28% were homemakers, 27% were private employers, 23% were self-employ, 18% were daily wages, and the remaining 3% had a government job. In the majority of 3rd years, students gained knowledge 58% from their teachers/tutors, 35 % from mass media and 7% from health articles.

Section-II

Assessment of knowledge on management of Hospital Acquired infections in children

It shows that all 60, 53(88.3 %) had inadequate knowledge, 7(11.7 %) had moderate knowledge and adequate knowledge was zero. Table.1 summarizes statistical outcomes of knowledge on preventing Hospital Acquired infections in children was 14.06 with SD 3.519 before S.T.P. The mean score percentage was computed, and it was 46.86%. The results found that the sampled subjects had inadequate knowledge regarding the management of Hospital Acquired infections in children. Thus from the statistical significance, it may be confirmed that knowledge of the management of Hospital Acquired infections in children increased after S.T.P.

Table 1: shows the knowledge score after a planned teaching program.

Knowledge	Mean	S. D	Mean %	't' value	Result
Pretest	14.06	3.519	46.86%	20.215	HS
Post -test	24.16	2.985	80.53%		P<0.05

Section-III

Evaluating the effectiveness of P.T.P. overall pretest and post-test mean knowledge on preventing Hospital Acquired infections in children. Table. 2 shows the pre and post-test knowledge of preventing Hospital Acquired infections.

Table 2: pre and post-test knowledge of management of Hospital Acquired infections.

Aspects	Max. score	Respondent's knowledge			Paired 't'-test
		Mean	SD	Mean (%)	
Pre-test	40	14.06	3.519	46.86%	20.215
Post-test	40	24.16	2.985	80.53%	

The study shows that the pretest findings depict that the pretest knowledge of preventing Hospital Acquired infections in children, such as incidence and cause aspect of management of Hospital Acquired infections, the mean score is 28.63%. Pattern and route of transmission of Hospital Acquired infections mean score is 28.33%, management of Hospital Acquired infections in children mean score is 37.60%. The mean combined score for preventing Hospital Acquired infections in children is 33.05% and SD is 3.519. In post-test knowledge of preventing Hospital Acquired infections in children, such as incidence and cause, the mean score is 52%. Pattern and route of transmission of Hospital Acquired infections mean score is 50.66%. Management of Hospital Acquired infections in children's mean score is 82.65%. and combined mean score of management of Hospital Acquired infections in children mean the score is 67.02%, and SD is 2.985. Improvement of knowledge regarding the management of Hospital Acquired infections in children, such as the general aspect of incidence and cause mean score, is 23.37%. Pattern and route of transmission mean score is 23.33%. Management of Hospital Acquired infections in children's mean score is 45.05%, the combined mean score of environmental hazards affecting the outcome of pregnancy mean score is 33.97%, and SD is 6.50. Paired 't' value of management of Hospital Acquired infections in children, combined paired 't'-test is 20.21.

Section-IV-Association of Post-test knowledge with selected demographic variables

Chi-square results of socio-demographic and post-test knowledge management of Hospital Acquired infections in children among B.Sc Nursing students are shown in Table.3.

Table 3: Chi-square results for sample (n=60)

Sl no	Characteristics	Chi-square Value	Results	P' value
1	Age	2.891	Non-Significant	>0.05
2	Gender	0.17	Non-Significant	<0.05
3	Religion	3.723	Non-significant	>0.05
4	Type of family	1.835	Non-significant	>0.05
5	Father's occupation	2.860	Non-Significant	>0.05

6	Source of information	6.25	Significant	<0.05
---	-----------------------	------	-------------	-------

Health can neither be demanded nor given. It can neither be bought nor sold, but the circumstances and services that are prerequisites to health can centrally be demanded and received as right. A protective environment in the hospital unit is essential, particularly when considering the services provided in the pediatric wards. Providing a safe and protective environment is a priority need. The present study was designed to assess the effectiveness of structured teaching program knowledge regarding preventing Hospital acquired infections in children among B.Sc Nursing students of VMP college of nursing at Akluj. The data findings have been organized and discussed based on the objectives and hypothesis.

The majority of students belong to the age group 21-22, i.e., 47% and 58% of students were female. 53% of students belonged to Christianity, and a maximum of 47% of them were from joint families. 28% of the sample's fathers were housemakers, and 58% of the samples were getting the source of information regarding the management of Hospital acquired infections from their teachers. The overall mean score percentage of knowledge is 46.86%; the current study investigator concludes that the selected student's knowledge regarding the management of Hospital Acquired infections in children was inadequate. The knowledge of students was influenced by the source of information. The calculated chi-square value for association of knowledge with the source of information was (Highly significant). The ability of students was also influenced by demographic variables such as a source of health information.

In the majority of B.Sc Nursing, students have inadequate knowledge about the incidence, causes, pattern of transmission and management of Hospital acquired infections in children. Education of students in these aspects is essential for the effective management of Hospital acquired infections in children. It aims to provide students with knowledge on identifying and avoiding trigger factors, maintaining asepsis and regular use of barrier techniques to safeguard children. Thus it may decrease the duration of hospital stay and provide comprehensive nursing care. It was found that the majority of B.Sc Nursing students' knowledge management of Hospital acquired infections in children was inadequate. It needs education in identifying and avoiding triggers,

sterilization and disinfection, and strict asepsis for proper preventive measures to be implemented in pediatric wards. The structured teaching program is beneficial to the nursing students in preventing Hospital acquired infections. This reduces mortality and morbidity associated with Hospital acquired infections.

Conclusions

The structured teaching program's efficiency in preventing Hospital Acquired infections in children has improved significantly among B.Sc Nursing students. The pretest revealed that B.Sc Nursing students understanding of Management of Hospital Acquired infections in children was lacking in all categories. It emphasizes the significance of regular in-service education sessions to keep students up to date on the latest developments in the management of Hospital Acquired infections in children. The post-test scores showed an improvement in knowledge after P.T.P. was administered. Hence it was concluded that Planned teaching program was an effective strategy for improving the knowledge of the B.Sc Nursing students.

Acknowledgment

The authors would like to express their gratitude to the Institutions of Nursing at Akluj for their help in conducting this study and the study participants for their whole hearted support.

Ethical clearance: Taken from institutional ethical committee.

Funding

None

Conflicts of Interest

None

References

1. Park K. Preventive and social medicine. Jabalpur.
2. Bitsori M, Maraki S, Koukouraki S, Galanakis E. Pseudomonas aeruginosa urinary tract infection in children: risk factors and outcomes. *The Journal of urology*. 2012 Jan;187(1):260-4.
3. Ducl G, Fabry J, Nicolle L. Prevention of hospital acquired infections: a practical guide. *Prevention of hospital acquired infections: a practical guide..* 2002(Ed. 2).
4. Coffin SE, Zaoutis TE. Infection control, hospital epidemiology, and patient safety. *Infectious Disease Clinics*. 2005 Sep 1;19(3):647-65.
5. Barr CE, Schulman K, Iacuzio D, Bradley JS. Effect of oseltamivir on the risk of pneumonia and use of health care services in children with clinically diagnosed influenza. *Current medical research and opinion*. 2007 Mar 1;23(3):523-31.
6. Garg BS, Chhabra S, Zothanzami SM. Safe motherhood: social, economic, and medical determinants of maternal mortality. *Women and Health Learning Package, The Network: TUFH Women and Health Taskforce*. Karachi: The Network. 2006 Sep;1132.
7. Romoşan O, Bocşan IS. The existing profile of nosocomial infections in municipal hospitals of general surgery. *Revista Medico-chirurgicala a Societatii de Medici si Naturalisti din Iasi*. 2011 Oct 1;115(4):1231-6.
8. El Ayyat AA, Sayed HA, Abou Had AM. A KAP study among staff and student nurses about infection control in Theodor Bilharz Hospital. *Journal of the Egyptian Society of Parasitology*. 2000 Aug 1;30(2):511-22.
9. Wheeler DS, Giaccone MJ, Hutchinson N, Haygood M, Bondurant P, Demmel K, Kotagal UR, Connelly B, Corcoran MS, Line K, Rich K. A hospital-wide quality-improvement collaborative to reduce catheter-associated bloodstream infections. *Pediatrics*. 2011 Oct;128(4):e995-1007.
10. Nichols RL. Preventing surgical site infections: a surgeon's perspective. *Emerging infectious diseases*. 2001 Mar;7(2):220.
11. Askarian M, Honarvar B, Tabatabaee HR, Assadian O. Knowledge, practice and attitude towards standard isolation precautions in Iranian medical students. *Journal of Hospital Infection*. 2004 Dec 1;58(4):292-6.
12. Klevens RM, Edwards JR, Richards Jr CL, Horan TC, Gaynes RP, Pollock DA, Cardo DM. Estimating health care-associated infections and deaths in US hospitals, 2002. *Public health reports*. 2007 Mar;122(2):160-6.
13. Nseir S, Grailles G, Soury-Lavergne A, Minacori F, Alves I, Durocher A. Accuracy of American Thoracic Society/Infectious Diseases Society of America criteria in predicting infection or colonization with multidrug-resistant bacteria at intensive-care unit admission. *Clinical microbiology and infection*. 2010 Jul 1;16(7):902-8.
14. Sydnor ER, Perl TM. Hospital epidemiology and infection control in acute-care settings. *Clinical microbiology reviews*. 2011 Jan;24(1):141-73.