

Study of the Orbital Vessels by Color Doppler in Known Diabetic Patients for Evaluation of Diabetic Retinopathy

Sushma Aharwal¹, Vishal Singh Rathore², Rekha Agrawal³, Susan Abraham⁴

¹Associate Professor Radiodiagnosis, Superspeciality Hospital NSCB Medical College Jabalpur (M.P.),

²Associate Professor Radiodiagnosis, ³Designated Professor Radiodiagnosis, N.S.C.B. Medical College Jabalpur (M.P.), ⁴Consultant Seth M. J. Hospital Jabalpur (M.P.).

How to cite this article: Sushma Aharwal, Vishal Singh Rathore, Rekha Agrawal et al. Study of the Orbital Vessels by Color Doppler in Known Diabetic Patients for Evaluation of Diabetic Retinopathy. Indian Journal of Public Health Research and Development 2023;14(2).

Abstract

Introduction: Diabetic retinopathy is a vascular disorder affecting the microvasculature of retina. It is caused by changes in the blood vessels of retina. If untreated, it may lead to blindness which is usually preventable if retinopathy is diagnosed early and treated promptly. As the prevalence of diabetes is rising, the systemic complications that include retinopathy, nephropathy, neuropathy and involvement of cardiovascular system are also increasing. Diabetic retinopathy is the leading cause of blindness in the world. Prevention of retinopathy needs early diagnosis. ⁽¹⁾ In ophthalmology, Color Doppler imaging is a new method that enables us to assess the orbital vasculature. It allows for simultaneous two dimensional anatomical and Doppler evaluations of retinal artery.

Objective and Aim: Aim of the study is to evaluate the ocular blood flow in patients with diabetes mellitus with no ocular symptoms.

Materials and Methods: Color Doppler evaluation of 50 diabetic patients (100 eyes) was done with SIEMENS SONOLINE G- 50 machine with High frequency probe (5-7.5 Mega Hertz). Doppler spectral analysis of ophthalmic arteries (OA) and central retinal arteries (CRA) were done. The peak systolic velocity (PSV), end diastolic velocity (EDV), resistive index (RI) and S/D ratio were calculated. PSV, EDV, RI were measured in all patients in both the eyes.

Results: The PSV of CRA in diabetics was significantly reduced ($p < 0.05$). The EDV of CRA in diabetics was also significantly reduced ($p < 0.05$). The RI of CRA is significantly increased ($p < 0.05$) in diabetics. The 95% confidence interval is observed in PSV, EDV, RI of CRA in diabetics.

Conclusion: There was statistical significant difference between the PSV, EDV and RI of CRA in diabetics. This significant difference could be due to the circulatory changes in blood vessels in diabetics. No significant difference was made in OA in diabetics. This study concludes that retinal hemodynamic changes were present even before the clinical manifestations of retinopathy in diabetics.

Keywords: Central retinal artery, ophthalmic artery, hemodynamic, resistance, diabetic.

Corresponding Author: Vishal Singh Rathore, Associate Professor, Radiodiagnosis Department, Netaji Subhash Chandra Bose Medical College Jabalpur (M.P.), India.

E-mail: drvishalrathore@gmail.com

Tel. 7982435897

Background

Despite the improvement in ophthalmoscopic examination in outpatient department (OPD) in Ophthalmology, in order to diagnose early changes in retinal arterial flow velocity, a newer imaging modality may be used for diagnosis of early changes in central retinal artery before clinical manifestation of retinopathy. Color Doppler imaging is the most promising modality that produces conventional gray-scale ultrasound images along with information regarding the direction and velocity of blood flow.⁽²⁾ The present study was attempted to evaluate clinically diagnosed diabetic patients without retinopathy by measuring the retinal arterial RI by duplex Color Doppler study. Retinal artery ultrasound, Color Doppler imaging was first used to image various organ systems in 1979. Later in 1989 Color Doppler imaging in orbit was described by Erickson. Eye is located superficially and cystic in nature. The normal anatomy and vasculature can be clearly seen by high frequency transducer. Color Doppler imaging of eye is a non-invasive procedure. It allows visualization of the grey scale imaging and color coded imaging both at the same time⁽³⁾. The peak systolic and end diastolic velocities of the ophthalmic and central retinal arteries can be measured using Doppler. The resistive index can then be calculated using peak systolic and end diastolic velocities. Orbital blood flow velocity can be qualitatively and quantitatively measured by color Doppler imaging.⁽⁴⁾ As Doppler shift detection sensitivity is higher than conventional grey scale resolution, evaluation of very small vessels supplying the orbit can be done non invasively. Wolfgang E. Lieb et al examined 40 normal eyes and they were able to locate the central retinal artery (CRA), posterior ciliary arteries (PCA) and ophthalmic arteries (OA) in all patients. Using Doppler spectrum, the blood flow velocity in these vessels are assessed quantitatively.

Anatomy Of Retina - Retina is a sensory tissue which lines the back of the eye. It is multilayered (10 layers) and contains photoreceptors namely rods and cones. The rods and cones convert light energy into signals which are then carried to brain through optic nerves and interprets the signal as visual images. Tiny blood vessels in retina take the oxygen and essential nutrients to the walls of the retina. In the centre of the

retina, there is a simple dimple called fovea, which is responsible for the sharp vision in eye. Optic nerve is a collection of nerve fibers which carries electric signals from retina to brain. Retina is supplied by central retinal artery which supplies the inner retinal layer and choroidal arteries which supplies the outer retinal layers. Central retinal artery is a branch of ophthalmic artery. Choroidal artery is a branch of posterior ciliary artery. Diabetic retinopathy is a vascular disorder affecting the microvasculature of retina. It is caused by changes in the blood vessels of retina. If untreated, it may lead to blindness. Therefore, if diagnosed and treated promptly, blindness is usually preventable.^(5,6) In ophthalmology, color Doppler imaging is a new method that enables us to assess the orbital vasculature. It allows for simultaneous two dimensional anatomical and Doppler evaluations of hemodynamic characteristics of retinal artery.^(7,8)

Material and Method

Color Doppler Imaging (CDI) is a safe, non invasive and highly reproducible procedure for evaluating hemodynamic alterations in the blood vessels. Color Doppler Imaging (CDI) combines with two dimensional (2D) ultrasonography and Doppler spectral analysis to evaluate the vascular structure. Blood flow velocities and flow pattern in orbital vessel are very useful for early detection of diabetic retinopathy.

Indices of Measurement

- (1) Resistance index (RI)
- (2) Peak systolic velocity (PSV)
- (3) End diastolic volume (EDV)

Methods

Retro bulbar blood flow velocity were assessed by using orbital Doppler and gray-scale sonography by SIEMENS SONOLINE G 50 Color Doppler machine. Imaging of the eyes were performed in all individuals by using a Color Doppler with a 5-7.5 MHz linear-array transducer. The patients were placed in the supine position and USG gel was applied to closed eyelids and 2D and color Doppler images were obtained by using high frequency linear probe. Ophthalmic artery was identified in the nasal

side of an eyeball, superior to the optic nerve, and it abuts the visible hypo reflective stripe representing the optic nerve. The central retinal artery arises from the ophthalmic artery and can be found anterior to the optic nerve, which is around 7.5 mm behind the ocular bulb. The posterior ciliary arteries are also supplied with blood by the ophthalmic artery, and they divide into multiple branches to supply the pial arteries. These arteries have a diameter of around 0.2 mm and form the pial network which adheres to the optic sheath and contributes to the vascularization of the optic nerve.


STUDY DESIGN- Prospective Study.

INCLUSION CRITERIA-


- Diabetic patients without retinopathy.
- Age group of 40-70 years.- Both sexes were included in the study.

EXCLUSION CRITERIA - Patients having infections or inflammatory lesions, benign or malignant lesions in orbit.


Statistical Analysis plan : All the records will be recorded by using structural schedule (Case Report Forms) and entered in Microsoft Excel Sheet. All the records will be rechecked for their completeness and consistencies. Non numeric entries will be coded numerically into nominal / ordinal distribution before analysis. Categorical variables was summarized in frequency and percent distribution and Chi-square or Fishers exact test will be performed as appropriate.



Color Doppler image showing high resistance flow in ophthalmic artery.



Color doppler image showing high resistance flow in Posterior ciliary artery.



Color Doppler evaluation of central retinal artery showing high resistance flow

Results

We studied 50 patients (100 globes), 30 men and 20 women, divided into age groups, average age was 55 years. Majority (54%) of subjects were in 5th decade of life with predominance of males. The mean duration of diabetes was 8.4 ± 1.2 years. Mean peak systolic velocity (PSV) in 50 diabetic patients without retinopathy was 8.90 ± 1.50 cm/sec ranging 5.30–11.01 cm/sec. Mean end diastolic velocity (EDV) in 50 diabetic patients without retinopathy was 3.21 ± 0.59 cm/sec ranging 2.5 –5.10 cm/sec . Mean resistive index (RI) in 50 diabetic patients without retinopathy was 0.92 ± 0.05 ranging 0.87–0.91.

Table 1: Distribution of patients according to age

Age	Number	Percent
40-50 Years	09	18%
51-60 Years	27	54%
61-70 Years	14	28%
TOTAL	50	100

Maximum number of patients seen in 51-60 (54%) years of age and minimum number of patients seen in 40-50 (18%) years of age.

Table 3: Average of PSV, EDV and RI in ocular vessels

VESSELS	PSV (Mean)	EDV (Mean)	RI (Mean)
Ophthalmic artery	8.11+ ₋ 1.12 cm/sec	3.4+ ₋ 1.2 cm/sec	0.57+ ₋ 0.08 cm/sec
Central retinal artery	8.90 + ₋ 1.50 cm/sec	3.21+ ₋ 0.59 cm/sec	0.92+ ₋ 0.05 cm/sec
Posterior cilliary artery	12.0+ ₋ 1.20 cm/sec	4.21+ ₋ 1.1 cm/sec	0.92+ ₋ 0.04 cm/sec

PSV, EDV and RI was increased in diabetic patients without retinopathy in all vessels.

Discussion

Due to characteristics eyeball location and constitution, ocular ultrasound has become an ideal method for the visualization of ocular structures and for the diagnosis of diseases that cannot be displayed on the fundus examination. The color Doppler ultrasound and pulsed Doppler has the advantage of being easily accessible, besides having no ionizing radiation and repeatedly performed as often as necessary, without leading to any additional risk to the patient, and do not need any medication prior to examination. Diabetes causes disturbances in the microcirculation by endothelial dysfunction⁽⁹⁾ causing perfusion disorders⁽¹⁰⁾, which will result in ultrasound significant decrease VPS in patients with ocular ischemic syndrome⁽¹¹⁾, central retinal artery occlusion⁽¹²⁾ and venous thrombosis⁽¹³⁾. The flow of the CRA should be antegrade, low resistance, with rounded systolic peak and continuous flow in diastole⁽¹⁴⁾. A study done by Mendivil A, Cuartero V, Mendivil MP,⁽¹⁵⁾ Titled -Ocular blood flow velocities in patients with proliferative diabetic retinopathy and healthy volunteers: a prospective study. British Journal of ophthalmology he compared 43 blood flow velocity in ocular vessels (ophthalmic artery, posterior ciliary arteries, central retinal vessels, and vortex veins) of 25 patients and showed that the diabetic patients had lower blood velocities than the volunteers. Ocular blood flow velocity was decreased

Table 2: Distribution of patients according to gender

Gender	Number	Percent
Male	30	60%
Female	20	40%
Total	50	100%

Maximum number of patients were male (60%), and female were (40%) in our study.

in diabetic patients with proliferative diabetic retinopathy. Schmetterer L, Wolzt M.⁽¹⁶⁾ who assessed Ocular blood flow and associated functional deviations in diabetic retinopathy. The total number of subjects included in this study is 80. Each group containing 40 patients. Both eyes were examined under this study. Hence there are a total of 160 eyes with 80 eyes in each group.

Conclusion

From the present study it can be concluded that, there is statistically significant increase in retinal arterial RI of type 2 diabetic patients without retinopathy. Increased RI in CRA, OA, and PCA showed that RI can be used to assess a index in the progression of diabetic retinopathy in patients or can also be used post pan retinal photocoagulation. Even though further investigations are needed to assess orbital hemodynamics in diabetic retinopathy, based on our study the results suggest that Color Doppler imaging has the ability to give the information on hemodynamic changes and can be used as a supportive modality for diagnosis of diabetic retinopathy in patients.

Conflict of interest – None to declare.

Acknowledgement – None to declare.

Ethical clearance – Taken by Ethical Committee

Source of Funding- None

References

1. Prahs P, Helbig H. Diabetic eye disease . The Umsch 2009 66: 183-188
2. Benden CJ, Abbitt PL, Beadless KA. Colour Doppler US of the orbit. Radio Graphic 1995; 15: 589-608
3. Gracner T. Ophthalmologica 2004 jul-aug;218 94); 237-42. Ocular blood flow velocity determined by colour Doppler imaging in diabetic retinopathy.
4. Gil Hernandez MA, Abreer Reyes P,Quintero M, Ayala E arch SOC ESP oftalmol. 2001 mar, 76 (3) 175 – 80 Doppler ultrasound in type 1 diabetes.
5. Sandeman DD, Shore AC, Tooke JE. Relation of skin capillary pressure in patients with insulin-dependent diabetes to complications and metabolic control. N Engl JMed 1992; 327: 760-764
6. Dandona R, Dandona L. Review findings from the Andhra Pradesh eye disease study. Indian J Ophthalmol 2001; 49: 215-234.
7. Mackinnon JR, McKillop G, Brien O, Butt Z, Nelson P. Colour Doppler imaging of the ocular circulation in diabetic retinopathy. Acta Ophthalmol 2000; 78: 386-389
8. Goebal W, Leib H, Sergott RC, Farhoumand R, Grehn F. Colour Doppler imaging: a new technique to assess orbital blood flow in patients with diabetic retinopathy. Invest Ophthalmol Vis C 1995; 36: 864-870.
9. Mendivil Antonio, Cuartero Victoria, Mendivil Maria, Ocular blood velocities in patients with proliferative diabetic retinopathy and healthy volunteers: a prospective study, British Journal of Ophthalmology 1995; 79; 413-416
10. GIL HERNÁNDEZ MA, ABREU REYES P,QUINTERO M, AYALA E. Ecografía Doppler en diabetes tipo I: primeros resultados Archivos de la sociedad española de oftalmología, Número 3. Marzo 2001
11. Ho AC, Lieb WE, Flaharty PM, Sergott RC, Brown GC, Bosley TM et al. Color Doppler imaging of the ocular ischemic syndrome. Ophthalmology 1992; 99: 1453-1462.
12. Sergott RC, Flaharty PM, Lieb WE et al. Color Doppler imaging identifies four syndromes of the retrobulbar circulation in patients with amaurosis fugax and central retinal artery occlusions. Trans Am Ophthalmol Soc 1992; 90: 383-401.
13. Keyser BJ, Flaharty PM, Sergott RC, Brown GC, Lieb WE, Annesley WH. Color Doppler imaging of arterial blood flow in central retinal vein occlusion. Ophthalmology 1994; 101:1357-1361.
14. Dr. José Herminio Vargas García, et al. Utilidad del ultrasonido Doppler orbitario en pacientes con enfermedades crónicas degenerativas. Presentación de hallazgos y revisión bibliográfica. Anales de Radiología México 2011;4:233-246.
15. Mendivil A, Cuartero V, Mendivil MP. Ocular blood flow velocities in patients with proliferative diabetic retinopathy and healthy volunteers: a 83 prospective study. British Journal of ophthalmology. 1995 May 1;79(5):413-6
16. Schmetterer L, Wolzt M. Ocular blood flow and associated functional deviations in diabetic retinopathy. Diabetologia. 1999 Mar 1;42(4):387-405.